

Geological mapping of an interesting lunar site: Tsiolkovski crater

G. Tognon¹, R. Pozzobon², M. Massironi²


¹ Center of Studies and Activities for Space “G. Colombo” (CISAS), UNIPD

² Department of Geosciences, UNIPD

© G. Tognon, R. Pozzobon & M. Massironi. All rights reserved.

Tsiolkovski crater


- Far side
- Feldspathic Highlands Terrane
- Oblique impact NW-SE
- Best example of farside mare volcanism (*Pieters & Tompkins, 1999*)
- Elevation floor difference ~450m (*Mouginis-Mark & Boyce, 2017*)
- Well-preserved central peak Ø
- Detections of OL and PAN



★ Olivine (*Corley et al., 2018*)

★ Purest Anorthosite (*Ohtake et al., 2009; Lemelin et al., 2015*)

20.4°S 129.1°E



© G. Tognon, R. Pozzobon & M. Massironi. All rights reserved.



Data

➤ TOPOGRAPHIC DATA:

- LRO LOLA and KAGUYA DEM Merge (*Barker et al. 2016*)
- horizontal resolution of 59 mpp
- vertical resolution of 3-4 m

$$S_m = R_r \times 2000$$

S_m = mapping scale
 R_r = raster resolution
 (Tobler, 1987)

➤ MONOCHROME MOSAICS:

- LRO WAC (*Robinson et al., 2010*) Global Mosaic
- resolution of 100 mpp
- LRO NAC (*Robinson et al., 2010*) mosaic
- resolution of 0,5 mpp (downscaled to 3 mpp)

Geomorphological mapping

$$S_m = 1:200.000$$

High-resolution
geological mapping

$$S_m = 1:6.000$$

➤ COLOR MOSAIC:

- CLEMENTINE UVVIS Color Ratio (*Lucey et al. 2000*)
- resolution of 200 mpp


R: 750/415 nm G: 750/950 nm B: 415/750 nm

Spectral mapping

$$S_m = 1:400.000$$

© G. Tognon, R. Pozzobon & M. Massironi. All rights reserved.

Geomorphological mapping



Geological contacts

- contact, approximate
- contact, certain

Morphologies

- crest of crater rim $D < 20$ km
- crest of crater rim $D > 20$ km
- mare rill

Structures


- wrinkle ridge

Geological units

- CF smooth material
- CF hummocky material
- CF central peak
- CW smooth ponds
- CW steep scarps
- CW rim

© G. Tognon, R. Pozzobon & M. Massironi. All rights reserved.

Spectral mapping



Spectral contacts


- contact, approximate
- contact, certain

Spectral units


- Fresher basalt
- Basaltic plain
- Noritic anorthosite - troctolite
- Anorthosite
- Mature highland soil

© G. Tognon, R. Pozzobon & M. Massironi. All rights reserved.

High-resolution geological mapping



$S_M = 1 : 6.000$!



Contacts



- contact, approximate
- contact, certain

Morphologies

- mare rill
- crater, slope $> 15^\circ$

Structures

- wrinkle ridge



© G. Tognon, R. Pozzobon & M. Massironi. All rights reserved.



Future works

- ✓ Finalization of the high-resolution geological mapping by means of the LRO NAC 3 mpp mosaic, implemented also to 0.5 mpp
- ✓ Radar investigation for the presence of deep structures, such as voids and lava pile emplacements, by means of Kaguya LRS (*Ono et al., 2010*) data
- ✓ Complete characterization (e.g. steepness analysis, boulders and craters counting for hazard analysis, landing ellipses and route traverses definition) as a possible landing site



Bibliography

- Pieters, C.M. & Tompkins, S., JGR, Vol. 104, pp. 21935-21949, 1999
- Mouginis-mark, P.J. & Boyce, J.M., Proceedings of the LPSC XLVIII, 2017
- Corley, L.M. et al., Icarus, Vol. 300, pp. 287-304, 2018
- Ohtake, M. et al., Nature, Vol. 461, pp. 236-241, 2009
- Lemelin, M. et al., JGR: Planets, Vol. 120, pp. 869-8878, 2015
- Barker, M.K., et al., Icarus, Vol. 273, pp. 346-355, 2016
- Robinson, M.S. et al., Space Sci. Rev., Vol. 150, pp. 81-124, 2010
- Lucey, P.G. et al., JGR, Vol. 105, pp. 20377-20386, 2000
- Tobler, W., Proceedings of the Land Resources Information Systems Conference, 12-16, 1987
- Ono, T. et al., Space Sci. Rev., Vol. 154, pp. 145-192, 2010

© G. Tognon, R. Pozzobon & M. Massironi. All rights reserved.