

# Latest Cretaceous foraminiferal ecology and palaeoceanographic inferences from chamber-specific LA-ICPMS analysis.

Michael J. Hennehan<sup>1,2</sup>, David Evans<sup>3</sup>, Wolfgang Müller<sup>3</sup>,  
Pincelli M. Hull<sup>2</sup>

<sup>1</sup>GFZ German Research Centre for Geosciences – Helmholtz Centre Potsdam, Telegrafenberg, 14473 Potsdam, Germany

<sup>2</sup>Department of Geology and Geophysics, Yale University, New Haven, Connecticut 06520, USA

<sup>3</sup>Institute for Geoscience, Goethe University Frankfurt, Frankfurt, Germany

# Motivations

- Recent work (e.g. Henehan et al. 2019, Woelders et al. 2018) has demonstrated the potential for application of boron isotope and trace element proxies to late Cretaceous foraminifera.
- However, to get accurate sea surface temperature and  $p\text{CO}_2$  estimates, we need to know which species to use, and what imprint their physiology may impart on their recorded proxy signals, as 'vital effects'.


# Motivations

- Numerous studies have used  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  to infer depth ranking, symbiosis and calcification pathways.
- However results are often contradictory:
  - relative depth habitats vary considerably within studies (e.g. Abramovich et al. 2003)
  - Inferences for symbiosis (e.g. *R. rugosa*, Abramovich et al. 2003) are often countered by other studies (e.g. D'Hondt and Zachos 1993, Falzoni et al. 2014)
- Since calibration of foraminiferal boron isotopes is key (e.g. Henehan et al. 2016, 2020), this makes CO<sub>2</sub> reconstruction more difficult.

# Approach


- LA-ICPMS analysis, chamber-by-chamber (see Evans & Müller (2018), Müller et al. (2009) for methods)
- Allows resolution of ecophysiology similar to serial test dissection
- Mg/Ca: potentially less susceptible to ontogenetic fractionation and diagenesis than  $\delta^{18}\text{O}$
- B/Ca: which in modern foraminifera is typically higher in symbiotic planktics and lower in symbiont-barren species
- Compare to solution measurements & measurements of  $\delta^{11}\text{B}$ .

# Representative Results: ODP Site 1049




Colour denotes species, lines and markers denote different tested individuals.  
Chamber No. 1 = earliest chamber analysed, Highest number = final chamber.  
Biserial (*Pseudotextularia elegans* & *Heterohelix globulosa*), multiserial (*Racemiguembelina fruticosa*) and trochospiral (*Contusotruncana contusa* & *Rugoglobigerina rugosa*) taxa

# *Racemiguembelina fructicosa*




- Clear depth migration in *R. fructicosa* with depth
- However magnitude of change and extreme temperatures from early chambers suggest a non-thermal influence on Mg/Ca

# *Racemiguembelina fructicosa*



- This species often thought to be shallowest dwelling and most clearly photosymbiotic (e.g. Houston et al. 1999, Isaza-Londoño et al. 2006).. Hence natural target for B isotope-pCO<sub>2</sub> reconstruction?

# *Racemiguembelina fructicosa*



- But solution MC-ICPMS shows  $\delta^{11}\text{B}$  decreases with size, counter to modern symbiont-bearers
- Also low in B/Ca
- And awkward to crush all chambers to remove clays!





Fig. from microtax.org



# B/Ca: Serials vs. Trochospirals



- Despite supposed lack of symbionts (e.g. Isaza-Londoño et al. 2006), *Rugoglobigerina rugosa* and *Contusotruncana contusa* are relatively enriched in Boron.

# Takeaways: *R. fructicosa*

- Despite published evidence from carbon isotopes for symbiosis in the photic zone, Mg/Ca (and  $\delta^{11}\text{B}$ ) suggests progressive migration to cooler, deeper waters with ontogeny (even as  $\delta^{13}\text{C}$  is known to increase; Houston et al. 1999).
- In agreement with dissection-derived  $\delta^{18}\text{O}$  (Houston et al. 1999)
- However, extremely high Mg/Ca in early chambers may indicate ontogenetic changes in biomineralization pathways of multiserial foraminifera that influence Mg incorporation.

# Takeaways: Trochospirals

- Very high B/Ca (even in an ocean of lower [B]; Lemarchand et al. 2002) may hint at more fundamental differences in biomineralization between major clades of Cretaceous foraminifera.
- Also suggests B/Ca is not necessarily a good indicator of photosymbiont association in Cretaceous planktic foraminifera.

# With thanks to

Sophia Riethmeister (U. Potsdam)

Brian Huber (Smithsonian)

Noah Planavsky (Yale)

Gavin Foster (U. Southampton) & Thomas Chalk (U.  
Southampton)

# References

- Abramovich, S., Keller, G., Stüben, D., Berner, Z., 2003. Characterization of late Campanian and Maastrichtian planktonic foraminiferal depth habitats and vital activities based on stable isotopes. *Palaeogeography, Palaeoclimatology, Palaeoecology* 202, 1–29. [https://doi.org/10.1016/S0031-0182\(03\)00572-8](https://doi.org/10.1016/S0031-0182(03)00572-8)
- D'Hondt, S., Zachos, J.C., 1993. On stable isotopic variation and earliest Paleocene planktonic foraminifera. *Paleoceanography* 8, 527–547. <https://doi.org/10.1029/93PA00952>
- Evans, D., Müller, W., 2018. Automated Extraction of a Five-Year LA-ICP-MS Trace Element Data Set of Ten Common Glass and Carbonate Reference Materials: Long-Term Data Quality, Optimisation and Laser Cell Homogeneity. *Geostandards and Geoanalytical Research* 42, 159–188. <https://doi.org/10.1111/ggr.12204>
- Falzoni, F., Petrizzo, M.R., Huber, B.T., MacLeod, K.G., 2014. Insights into the meridional ornamentation of the planktonic foraminiferal genus *Rugoglobigerina* (Late Cretaceous) and implications for taxonomy. *Cretaceous Research* 47, 87–104. <https://doi.org/10.1016/j.cretres.2013.11.001>
- Henehan, M.J., Edgar, K.M., Foster, G.L., Penman, D.E., Hull, P.M., Greenop, R., Anagnostou, E., Pearson, P.N., 2020. Revisiting the Middle Eocene Climatic Optimum 'Carbon Cycle Conundrum' with new estimates of atmospheric  $pCO_2$  from boron isotopes. *Paleoceanography and Paleoclimatology*. <https://doi.org/10.1029/2019PA003713>
- Henehan, M.J., Foster, G.L., Bostock, H.C., Greenop, R., Marshall, B.J., Wilson, P.A., 2016. A new boron isotope-pH calibration for *Orbulina universa*, with implications for understanding and accounting for 'vital effects.' *Earth and Planetary Science Letters* 454, 282–292. <https://doi.org/10.1016/j.epsl.2016.09.024>
- Henehan, M.J., Ridgwell, A.J., Thomas, E., Zhang, S., Alegret, L., Schmidt, D.N., Rae, J.W.B., Witts, J.D., Landman, N.H., Greene, S.E., Huber, B.T., Super, J.R., Planavsky, N.J., Hull, P.M., 2019. Rapid ocean acidification and protracted Earth System recovery followed the end-Cretaceous Chicxulub impact. *Proceedings of the National Academy of Sciences* 116, 22500–22504. <https://doi.org/10.1073/pnas.1905989116>
- Houston, R.M., Huber, B.T., Spero, H.J., 1999. Size-related isotopic trends in some Maastrichtian planktic foraminifera: methodological comparisons, intraspecific variability, and evidence for photosymbiosis. *Marine Micropaleontology* 36, 169–188. [https://doi.org/10.1016/S0377-8398\(99\)00007-9](https://doi.org/10.1016/S0377-8398(99)00007-9)
- Isaza-Londoño, C., MacLeod, K.G., Huber, B.T., 2006. Maastrichtian North Atlantic warming, increasing stratification, and foraminiferal paleobiology at three timescales: NORTH ATLANTIC MAASTRICHTIAN WARMING. *Paleoceanography* 21, n/a–n/a. <https://doi.org/10.1029/2004PA001130>
- Lemarchand, D., Gaillardet, J., Lewin, É., Allègre, C.J., 2002. Boron isotope systematics in large rivers: implications for the marine boron budget and paleo-pH reconstruction over the Cenozoic. *Chemical Geology* 190, 123–140. [https://doi.org/10.1016/S0009-2541\(02\)00114-6](https://doi.org/10.1016/S0009-2541(02)00114-6)
- Müller, W., Shelley, M., Miller, P., Broude, S., 2009. Initial performance metrics of a new custom-designed ArF excimer LA-ICPMS system coupled to a two-volume laser-ablation cell. *J. Anal. At. Spectrom.* 24, 209–214. <https://doi.org/10.1039/B805995K>
- Woelders, L., Vellekoop, J., Weltje, G.J., de Nooijer, L., Reichert, G.-J., Peterse, F., Claeys, P., Speijer, R.P., 2018. Robust multi-proxy data integration, using late Cretaceous