

# Influence of grass invasion on soil parameters in a Belgian heathland

Natascha Arnauts<sup>1</sup>, Miguel Portillo Estrada<sup>2</sup>, Monique Carnol<sup>3</sup>, Bernard Bosman<sup>3</sup>, François Rineau<sup>1</sup>

<sup>1</sup>Centre for Environmental Sciences, Hasselt University, Diepenbeek, Belgium. <sup>2</sup>Centre of excellence (PLECO), University of Antwerp, Wilrijk, Belgium. <sup>3</sup>InBios, Plant and Microbial Ecology, University of Liège, Liège, Belgium



## Introduction

- N deposition
- Heather thrives on nutrient poor soils → now outcompeted by subdominant species: grass, better suited to these elevated N levels
- Heather (vs grass): woody litter → lignin → harder to degrade → slows down N cycle

**Hypothesis:** lower nutrient turnover in heathland compared to grassland → impact on soil nutrient cycling



Figure 1. Grass invaded heathland

## Method

- 14 Plots
  - grass gradient

- Measurement of:
  - Vegetation cover
  - Root biomass
  - Soil water content
  - N net mineralization rate (NMR)
  - Relative nitrification
  - Enzymes: chitinase & phosphatase
  - Total C & total N




Figure 2. Vegetation cover per plot

## Results




Figure 3. Effect of vegetation on N turnover

- No significant correlation  
→ vegetation has no impact on measured parameters
- Nonlinear trend: high NMR at threshold value of approximately 70% grass  
→ effect when vegetation is truly dominant




Figure 4. Correlation of phosphatase & chitinase

The data show significant correlations between **Chitinase & phosphatase**  
→ Tightly coupled N & P cycles




Figure 5. Correlation total C & net mineralization rate

### Total C & NMR

→ NMR = conversion of organic matter to NH<sub>4</sub>  
→ OM ↑ → NMR ↑ → C ↑




Figure 6. PCA of variables in function of different plots with grass gradient

2 components represents 85% of the total variation in the data:  
Component 1: 62,2%  
Component 2: 23,8%

Organic matter & NMR contribute most to dataset (positive association)

Most points=clustered, outliers: sample 8A, 10A, 11A, 11B, 4C, 7A  
(Total samples n=56)

## Conclusions

- The study shows
  - no impact of grass invasion on these parameters measured in the heathland
  - organic matter & NMR show an influence on the data
  - dominant vegetation → effect of litter
- Further research is needed to understand how to maintain ecosystem

## Acknowledgements

The authors would like to thank ANB (Agentschap voor Natuur en Bos) and NPHK (Nationaal Park Hoge Kempen) for allowing us to perform our experiments. This research is financed by BOF Doctoral Funds Hasselt University.

A special thanks to Laure Steenaerts, Wouter Reyns, Lennert Indestege, Jolien Wevers, Laura Coninx, Michiel Huybrechts and Jana Deckers for their help.