



Imaging the Earth's magnetic environment in soft X-rays with SMILE


Graziella Branduardi-Raymont¹, Steve Sembay², Tianran Sun³, Hyunju Connor⁴, Andrey Samsonov¹

¹ Mullard Space Science Laboratory, University College London, Holmbury St Mary, Dorking, UK


² University of Leicester, Leicester, UK

³ National Space Science Center, Chinese Academy of Sciences, Beijing, China

⁴ University of Alaska, Fairbanks, USA



EGU2020 “Sharing Geoscience Online” – Session ST2.5




SMILE




Solar wind Magnetosphere Ionosphere Link Explorer

SMILE Science

Investigate the dynamic response of the Earth's magnetosphere to the solar wind impact in a unique and global manner






SMILE mission and instruments




- SMILE is a joint scientific mission, from inception to launch and operations, by the **European Space Agency** and the **Chinese Academy of Sciences**, it is under development and due for launch at the end of 2023
- SMILE will investigate the dynamic response of geospace to the solar wind impact, exploring the **full chain of events that drive Space Weather**
- SMILE combines **X-ray imaging** of the dayside magnetosheath and the cusps (with the Soft X-ray Imager, **SXI**), simultaneous **UV imaging** of the Northern aurora (UltraViolet Imager, **UVI**) and in situ monitoring of the **solar wind** and **magnetosheath** conditions (Light Ion Analyser, **LIA**, and **MAG**netometer) from a very elliptical orbit




SMILE scientific motivations

Study the full chain of events that drive Sun-Earth relations




- Structure and dynamics of the magnetosphere mainly controlled by **magnetic reconnection**: Basic theory of magnetospheric circulation well known, **microscale** explored by many in situ measurements
- Reality of how complex interaction takes place on **a global scale**, and **how it evolves**, still not understood
- SMILE can answer questions which help **distinguishing modes of interaction**



SMILE scientific motivations

What are the fundamental modes of the dayside solar wind / magnetosphere interaction?


- When/where is **reconnection** steady/transient/bursty, patchy or global?
- Dependent on solar wind parameters or intrinsic instabilities?
- Component or anti-parallel
- Role of the magnetospheric cusps in solar wind/magnetosphere coupling



Ion energy decreases towards pole for IMF S, and vice versa

Cusps expands poleward after IMF turns N

Escoubet et al. 2008



SMILE scientific motivations

What defines the substorm cycle?


- Auroral oval responds to changes in magnetospheric or solar wind conditions: IMF orientation, dynamic pressure triggers?
- Other modes of magnetospheric behaviour: e.g. saw-tooth events, auroral beads


How do CME-driven storms arise?

How do they relate to substorms?




- Fast solar wind and long intervals of S IMF: Is solar wind driving the only storm trigger?

- Relation storm – substorm? How do storms end? Space weather relevance





SMILE Soft X-ray Imager (SXI)



Lobster-eye Micropore Optic

Ultra-wide field of view
~16° x 26°

Focal length 30 cm

Optic Mass < 1kg


Instrument ~36 kg

CCD Detector Plane


Photon counting: Photon lists with ~4 s time resolution

High QE in soft X-rays
~90% at 500 eV

Medium energy resolution
~50 eV FWHM at 500 eV



PI S. Sembay, Univ. of Leicester, UK



SXI performance versus requirements


The SMILE Consortium comprises a number of Working Groups whose aim is to help ensuring that the mission science objectives are achieved and optimised

Among these the **Modelling WG** dedicates its efforts to verifying that the SXI performance satisfies the scientific requirements of the mission

Two main requirements on SXI can be expressed as:


- For solar wind flux $> 4.9 \times 10^8 \text{ cm}^{-2} \text{ s}^{-1}$ the location of the subsolar magnetopause shall be determined with a better accuracy than $0.5 R_E$ and better than 5 min time resolution from locations on orbit greater than $15 R_E$ geocentric
- The poleward and/or equatorward edges of the mid-altitude cusp in X-ray images shall be identified with $0.25 R_E$ accuracy at a time resolution of at least 5 min for solar wind flux $> 4.9 \times 10^8 \text{ cm}^{-2} \text{ s}^{-1}$.

Solar Wind Charge eXchange (SWCX)



$$P_X = \alpha n_{sw} n_n \langle g \rangle \text{ eV cm}^{-3} \text{ s}^{-1}$$

X-ray emission proportional to density of solar wind ions and neutrals, hence brightest in the dayside magnetosheath and the cusps




17th March 2015 storm event

Before storm: 04 UT, $N=15 \text{ cm}^{-3}$, $V= 410 \text{ km/s}$

Solar wind input

X-ray emissivity line-of-sight integrated




**Predicted SWCX X-ray emissivity
from MHD simulation**


During storm: 04:25 UT, $N=50 \text{ cm}^{-3}$, $V= 510 \text{ km/s}$

Solar wind input

X-ray emissivity I-o-s integrated



T. Sun, NSSC, CAS, China






From X-ray emissivity to observed counts


Detectable X-ray emission
calculated by integrating
along the line of sight
through the modelled
X-ray emissivity cube

→ SXI_SIM produces expected count maps

140912_IMF_turning_larger_alpha_o3_1905
 N_{sw} : 22.69 cm⁻³ V_{sw} : 623.14 km s⁻¹ B_y : 7.89 nT B_z : 14.87 nT
 Position: 8.58 5.16 17.03 GSE
 Aim Point: 8.48 0.00 0.00 GSE




→ SXI performance study versus
SMILE science requirements



Extracting magnetopause location from SXI images: techniques under study (1)

Gradient method



Take locations along image slices on the Earth-Sun line through MHD simulated X-ray emissivity (left) and SXI count images (right), where


← X-ray emission gradient is greatest


↓ X-ray emission is greatest

Maximum method

True magnetopause location likely to be in between the two estimates

Credit: A. Samsonov








Extracting magnetopause location from SXI images: techniques under study (2)

Tangential direction approach

- The direction with maximum X-ray intensity is the tangential direction
- Use two-point observations to pinpoint the tangent point
- Position of the magnetopause is at the tangent point



Credit: : H. Connor & M. Collier



Extracting magnetopause location from SXI images: techniques under study (3a)


Boundary fitting in 3D

Reconstruction of 3D boundaries from 2D images:

Assume an X-ray emission model, and a boundary model extension in 3D

→ Model an ensemble of X-ray images

→ Fit to the observed image, best match gives position of magnetopause in 3D



Credit: A. Jorgensen & T. Sun


Emissions model (empirical)

$$F(\vec{r}) = \begin{cases} 0 & \text{inside MP} \\ (A_1 + B \sin^8 \theta) \left(\frac{r}{r_{\text{ref}}} \right)^{-(\alpha + \beta \sin^2 \theta)} & \text{between MP and BS} \\ A_2 \left(\frac{r}{r_{\text{ref}}} \right)^{-3} & \text{outside BS} \end{cases}$$

Boundary model (a natural extension of Shue et al. [1997])

$$r(\theta, \phi) = \frac{r_y(\theta) r_z(\theta)}{\sqrt{[r_z(\theta) \cos \phi]^2 + [r_y(\theta) \sin \phi]^2}}$$

$$r_y(\theta) = r_0 \left(\frac{2}{1 + \cos \theta} \right)^{\alpha_y} \quad r_z(\theta) = r_0 \left(\frac{2}{1 + \cos \theta} \right)^{\alpha_z}$$



Extracting magnetopause location from SXI images: techniques under study (3b)

Forward modelling


Application of Jorgensen & T. Sun technique to MHD simulations of SXI images


Model compared to dataset
via chi-square test

Parameters of empirical model
adjusted to minimise chi-square

Requirement of $0.5 R_E$ accuracy
on magnetopause location for
5 min exposure **comfortably**
satisfied

Credit: S Sembay






Extracting magnetopause location from SXI images: techniques under study (4)

Tangent fitting approach


Combines techniques (2) and (3):

- Assume a model for the magnetopause
- For each left free parameter calculate the tangent direction along all longitudes
- Find the best match
- Technique saves time, and avoids introducing models for X-ray emissivity and bow shock position




X-ray intensity inside FOV ($16^\circ \times 27^\circ$)
 Black curve: Position of the maximum X-ray intensity
 Red curve: Best fit tangential direction

Credit: T. Sun et al.



Extracting magnetopause location from SXI images: techniques under study (5)

Tomographic reconstruction





Reconstruct a 3D object from multiple 2D projections

Most significant limitation on reconstruction is probably the small range of viewing angles

Credit: A. Jorgensen & T. Sun



Images from typical orbit





Locating the magnetospheric cusps (1)



The point with apparent increase of **X-ray gradient** in the image corresponds well to the **tangential direction of the cusp boundary**




Red dots: From X-ray image, with sudden increase of the X-ray gradient

Black dots: Tangent direction, from MHD cusp boundary


Good agreement with each other. Average deviation is ~0.18 deg.



Credit: T. Sun




Locating the magnetospheric cusps (2)




Taking a slice through the MHD simulation

Credit: S. Sembay



Images on the right:

- View of the North (top) and South (bottom) cusps
- MHD simulation (left), SXI simulation (right)
- **Cusp edges can be determined with $0.25 R_E$ accuracy over 5 min as required**



Conclusions

SMILE aims to investigate the dynamic impact of the solar wind on the Earth's magnetosphere in a **global and **novel** way**

SMILE will provide direct **scientific input** to the studies of space weather by making the remote sensing measurements needed to **validate global models** of solar wind-magnetosphere interactions

In preparation of SMILE launch and operations:

- MHD simulations employed to produce expected SXI images of magnetosheath and cusps for different solar wind conditions
- Simulated SXI images used to establish the instrument performance by:



Applying a variety of techniques aiming to determine the location of the magnetopause

Extracting magnetospheric cusps location and extent

Results found to be consistent with SMILE scientific requirements



Thank you!



<http://sci.esa.int/smile/>

<http://english.cssar.cas.cn/smile/>

<http://www.mssl.ucl.ac.uk/SMILE/>