

Resolution dependence of magnetosheath waves in global hybrid-Vlasov simulations

Annales Geophysicae Discussions, angeo-2020-24

Maxime Dubart¹, Urs Ganse¹, Adnane Osmane¹, Andreas Johlander¹, Markus Battarbee¹, Maxime Grandin¹, Yann Pfau-Kempf¹, Lucile Turc¹ and Minna Palmroth^{1,2}

¹Department of Physics, University of Helsinki, Helsinki, Finland

²Space and Earth Observation Centre, Finnish Meteorological Institute, Helsinki, Finland



Simulations set-up

- Global hybrid-Vlasov model: VLASIATOR [1]
- Cartesian 2D spatial grid
- Cartesian 3D velocity grid
- Protons described as velocity distribution functions (VDFs)
- Electrons as a cold massless charge neutralizing fluid
- Closure of the system: generalised Ohm's law with Hall term
- Noon-midnight meridional plane (X-Z)
- Inner boundary at 4.7 RE from the centre of the Earth
- Solar wind: flowing in -X direction, 750 km/s
- IMF: 5 nT, 45 degrees angle with respect to X, southward
- Temperature: 0.5 MK
- Density: 1 cm⁻³

Three different spatial resolutions


$\Delta r = 300 \text{ km} = 0.76 \text{ di}$

$\Delta r = 600 \text{ km} = 0.38 \text{ di}$

$\Delta r = 900 \text{ km} = 0.25 \text{ di}$


Ion-scale waves

• Fast Fourier Transform




Left-handed → **Alfvén Ion Cyclotron waves**
circularly polarised **Mirror Modes [2]**


• Minimum variance analysis



• Growth rate



Velocity Distribution Functions




• Proton cyclotron and Mirror instabilities well resolved at $\Delta r = 300 \text{ km}$. Sufficient maximum resolution

• Proton cyclotron instability not resolved at $\Delta r = 900 \text{ km}$

• Temperature anisotropy increases at low resolution due to not resolving AIC waves

Temperature anisotropy



• $\Delta r = 440 \text{ km} = 0.6 \text{ di}$ presents an acceptable minimum resolution for study of magnetosheath waves in global hybrid-Vlasov

[1] Palmroth, M. et al., Vlasov methods in space physics and astrophysics, LRCA, 4, <https://doi.org/10.1007/s41115-018-0003-2>, 2018
[2] Hoilijoki, S. et al., Mirror modes in the Earth's magnetosheath: Results from a global hybrid-Vlasov simulation, JGR, 121, <https://doi.org/10.1002/2015JA022026>, 2016.

We acknowledge The European Research Council for Starting grant 200141-QuESpace, with which Vlasior was developed, and Consolidator grant 682068-PRESTISSIMO awarded to further develop Vlasior and use it for scientific investigations. The Finnish Centre of Excellence in Research of Sustainable Space (FORESAIL), funded through the Academy of Finland grant number 312351, supports Vlasior development and science as well. The CSC - IT Center for Science in Finland is acknowledged for the Sisu and Taito supercomputer usage, which led to the results presented here. Maxime Dubart thanks Thiago Brito for useful discussions throughout the study which helped carrying it out.