

# **How tail reconnection affects the asymmetric state of the magnetosphere in the LFM model**


**A. Ohma, N. Østgaard, J. P. Reistad, K. M. Laundal, P. Tenfjord**

**EGU 2020**

# IMF $B_y$ induces north-south asymmetries

IMF  $B_y$  is a source of numerous asymmetric features in our magnetospheric system, e.g. north-south asymmetries in the aurora, the magnetospheric and ionospheric current systems and the plasma convection.


Field-aligned and divergence free currents from the AMPS model with IMF  $B_y = -8$  nT



AMPS model: *Laundal et al. (2018)*

Asymmetrically displaced substorm onset for IMF  $B_y < 0$


November 15, 2001



*Østgaard et al. (2004)*

# How the asymmetry is induced

Dayside reconnection geometry when IMF  $B_y > 0$




Tenfjord et al. (2015)

Magnetic flux from dayside reconnection is added asymmetrically to the two hemispheres due to magnetic tension  
For IMF  $B_y > 0$ , the flux is added at dawn in NH and at dusk in SH

The resulting asymmetric pressure distribution induces asymmetries in the magnetosphere (*Khurana et al.*, 1996; *Liou & Newell*, 2010; *Tenfjord et al.*, 2015)

Closed nightside fieldlines at  $L = 11 R_E$ , 25 minutes after IMF  $B_y$  were introduced in the MHD simulation



Tenfjord et al. (2015)

# Timescales at geosynchronous orbit

Response to IMF  $B_y$  polarity changes at geosynchronous orbit  
(*Tenfjord et al.*, 2017, 2018)


Response time < 15 min

Reconfiguration time < 50 min

Similar timescales

- For both IMF  $B_y$  polarity changes
- At the dayside and at the nightside
- For IMF  $B_z < 0$  and  $B_z > 0$

Confirm that asymmetries in the lobe pressure play a major role in inducing these asymmetries.







# Reduced asymmetry during substorms

How do increased tail reconnection affect asymmetries?


*Østgaard et al. (2011, 2018) and Ohma et al. (2018):*

Reduced asymmetry during substorms

IMF  $B_y < 0$



Input IMF parameters



Run number at CCMC  
Anders\_Ohma\_082219\_1

To further investigate how enhanced tail reconnection affects the asymmetric state of the magnetosphere, we apply a global MHD model.

We use the LFM model, as LFM results most closely resemble “idealized substorm” (Gordeev et al., 2017)

Constant IMF  $B_y = -5$  nT

IMF  $|B_z| = 5$  nT


0 - 59 minutes: Southward IMF

60 - 179 minutes: Northward IMF

180 - 300 minutes: Southward IMF

The north-to-south polarity change “triggers” a loading-unloading cycle

# MHD model - results



**Closed field lines at the  
nightside are more symmetric  
after the increase in tail  
reconnection**

Consider the time interval 180 - 300 minutes

The amount of open flux increases in the interval 180 - 239 min ("growth" phase)  
"Onset" at  $t = 239$  min


The amount of open flux decreases in the interval 239 - 260 min ("Expansion" phase)  
The CTP also maximises in the period 239 - 260 min, implying enhanced tail reconnection  
The magnetic field lines are more symmetric after the unloading have occurred

Cross-tail potential (CTP) quantifies  
earthward flux transport in the  
closed magnetotail.


We have used:

$$CTP = - \sum_{-15}^{+15} (\mathbf{v} \times \mathbf{B})_y \Delta y$$

at  $X = -12 R_E$



# MHD model - evolution at a fixed location



Mapping the field line located at  $X = -12 R_E$ ,  $Y = 0 R_E$ ,  $Z = 0 R_E$  to both hemispheres for every second time step.

Most asymmetric in the time interval 208 - 230 min, then reduced asymmetry

Near symmetric for  $> 250$  min



# MHD model - a closer look




Figure of CTP shows several regions with enhanced earthward transport of flux, implying enhanced tail reconnection

$t = 231$ : Increase at dawn


$t = 242$ : Increase at dusk

Figures of longitudinal displacement show large displacement in all MLT sectors before nightside reconnection sets in.


Reduced asymmetry at dawn following the increase at  $t = 131$   
Reduced asymmetry at dusk following the increase at  $t = 242$

Consistent with the auroral observations, reduced asymmetry when tail reconnection increases.

Grid mapped from SH to NH to yield displacement



# MHD model - cross-section of the magnetotail



Cross-section of the magnetosphere showing magnetic pressure and plasma convection.

Blue contour indicate the magnetopause using the technique outlined in *Peng et al. (2010)*

Red contour indicate the open-closed boundary, determined by field line tracing


General increase in lobe pressure until  $t = 240$

Local decrease at dawn when tail reconnection starts here ( $t = 235$  and  $t = 240$ )

Global decrease when tail reconnection starts at dusk ( $t = 245$  and  $t = 250$ )

# Summary

- LFM run with constant IMF  $B_y = -5$  nT
- Following the southward turning of the IMF, the lobe pressure and the north-south asymmetry of the field lines increase
- The simulation clearly shows reduced asymmetry when tail reconnection sets in
- The reduction occurs first at dawn, then at dusk, which is consistent with the location of the reconnection region in the tail
- The regions of reduction also coincide with regions where the lobe flux is reduced
- The model is consistent with auroral observations, which show reduced asymmetry during substorms
- The reduced asymmetry is likely a result of changes in the force balance in the tail caused by the commencement of tail reconnection, and possibly directly related to the decrease of lobe pressure





- Gordeev et al., (2017)**, The substorm cycle as reproduced by global MHD models, doi:10.1002/2016SW001495.
- Khurana et al., (1996)**, Magnetospheric convection in the presence of interplanetary magnetic field  $B_y$ : A conceptual model and simulations, doi:10.1029/95JA03673
- Laundal et al., (2018)**, Solar Wind and Seasonal Influence on Ionospheric Currents From Swarm and CHAMP Measurements, doi:10.1029/2018JA025387
- Liou & Newell (2010)**, On the azimuthal location of auroral breakup: Hemispheric asymmetry, doi:10.1029/2010GL045537
- Ohma et al., (2018)**, Evolution of asymmetrically displaced footpoints during substorms, doi:10.1029/2018JA025869
- Peng et al., (2010)**, Role of IMF  $B_x$  in the solar wind-magnetosphere-ionosphere coupling, doi:10.1029/2010JA015454.
- Østgaard et al., (2018)**, Interplanetary magnetic field control of the location of substorm onset and auroral features in the conjugate hemispheres, doi:10.1029/2003JA010370
- Østgaard et al., (2011)**, Evolution of auroral asymmetries in the conjugate hemispheres during two substorms, doi:10.1029/2010GL046057
- Østgaard et al., (2018)**, The asymmetric geospacer as displayed during the geomagnetic storm on 17 august 2001, doi:10.5194/angeo-36-1577-2018
- Tenfjord et al., (2015)**, How the IMF  $B_y$  induces a  $B_y$  component in the closed magnetosphere and how it leads to asymmetric currents and convection patterns in the two hemispheres, doi: 10.1002/2015JA021579
- Tenfjord et al., (2017)**, Magnetospheric response and reconfiguration times following IMF  $B_y$  reversals, doi:10.1002/2016JA023018
- Tenfjord et al. (2018)**, How the IMF  $B_y$  Induces a Local  $B_y$  Component During Northward IMF  $B_z$  and Characteristic Timescales, doi:10.1002/2018JA025186