

QUANTITATIVE STRUCTURAL ANALYSIS OF FRACTURES USING DIGITAL OUTCROP MODELS

Caroline Modica Custódio²

Maria Alcione Lima Celestino⁴

Laís Vieira de Souza²

Jaqueline Lopes Diniz^{1,2}

Leonardo Campos Inocencio^{2,3}

Juliano Bonato^{1,2}

Aline Fernanda Spaniol²

Tiago Siqueira de Miranda^{2,4}

Francisco Manoel Wohnrath Tognoli^{1,2,*}

¹Graduate Program on Geology, Unisinos University, São Leopoldo, RS, Brazil

²VizGEO Research Group, Unisinos University, São Leopoldo, RS, Brazil


³Graduate Program on Applied Computing, Unisinos University, São Leopoldo, RS, Brazil

⁴Department of Geology, Federal University of Pernambuco, Recife, PE, Brazil

* ftognoli@unisinos.br


https://www.researchgate.net/profile/Francisco_Tognoli





Study area

- Location of the Araripe Basin in Northeastern Brazil
- Intracontinental rift basin positioned between the Patos and Pernambuco shear zones (Brazilian/Pan-African orogeny)



- Two different DOMs in this work (ARN02 & ARN04)
- They are on the fault zone context and represent the Araripe's Basin basement

ARN02

- Natural outcrop along a creek
- Approximately 35m long and 5m height
- Tilted metamorphic rocks
- Intensely fractured with breccia zones (see dashed lines)



ARN04

- Road pavement pit
- Approximately 45m long and 4m height
- Orthogneisses and amphibolites
- Intensely fractured (joints and faults), mostly perpendicular to the outcrop wall.



Field Data Collection
→ ARN02 & ARN04



Photographs captured directly from the ground




- GNSS RTK positioning system + ground control points
- Positional errors smaller than 2 cm

3D Model reconstruction

- Photogrammetric models
- Structure from Motion (SfM) technique
- Spatial resolution 0.3 mm/pixel

Example of SfM Photogrammetry workflow

Javadnejad (2018)



DOMs analysis


- Extract fracture patterns with CloudCompare
- Plugins:
 - Compass
 - Facets

Results & Interpretation

FACETS PLUGIN → PLANES EXTRACTION METHODOLOGY


- FACETS → plugin within CloudCompare (Dewez et al. 2016)
- Perform automatic planar facet extraction
- Segmenting massive 3D point clouds into individual planar facets
- Calculate their dip and dip direction
- Report the extracted data in interactive stereograms

Geological planar facets extracted are easily separated in families, and subfamilies sets → spatial orientation similarity (azimuth).




COMPASS PLUGIN → PLANES EXTRACTION METHODOLOGY

- COMPASS → plugin within CloudCompare (Thiele et al. 2017)
- Rapidly interpolate structural features between manually defined control points → point cloud and raster datasets
- Tools for measuring surface orientations, lineations and true thicknesses
- Map Mode → delineating geological units
- Compass Mode → measuring orientations and thicknesses
 - ↳ Plane tool → Measure surfaces orientation
 - ↳ Trace tool → Digitize and measure traces and contacts










Automatic planar extraction → FACETS SEGMENTATION

Original Point cloud




Point cloud manual clean-up




- Why is the point cloud manual clean-up an essential step before extracting the planar structures?
Reduce noise and point cloud size → optimize processing
- This step significantly increases the measurement accuracy (True planes x False planes)


RESULTS - ARN02 vs. ARN04





Data accuracy → 62%.




Data accuracy → 10,9%.



- Facets Tool in the ARN02 area: **EFFECTIVE**
- Outcrop wall orientation:
 - bedding planes (So) (green): perpendicular → large area
 - breccia zone (blue): fracture dip direction perpendicular and parallel
- Many true planes extracted (almost all the outcrop) → Why?
 - True planes are visible




- Facets Tool in the ARN04 area: **INEFFECTIVE**
- Outcrop wall orientation:
 - fracture dip direction parallel
- Few true planes extracted → Why?
 - less visible planes exposed to calculate automatically.
 - most of the fractures are exposed as lines (trace)



In cases like this (only line visible), manual planar extraction using the Compass tool is more effective.

COMPASS vs. Field Measurements - ARN04

- Structural Analysis comparing the DOM measurements (Compass - manual planar extraction tool) and field measurements




- **Field Measurements** (below the dotted line)

- 80 planes measurements
- 23 principal fractures

- **DOM Measurements** (overall outcrop)

- 86 planes measurements
- 23 principal fractures (same structures that were identified in the field)

- Fractures preferential orientation: $N45-50^{\circ} E$
- Dip: variation along the structure
- The fractures measurements in this outcrop are part of a negative flower structure, typical of the shear zone, generated by strike-slip tectonic efforts.




Negative Flower

FACETS vs. COMPASS

FACETS	COMPASS
Good tool for large scale surveys	Good tool for small scale surveys
Semi-automatic	Manual
Exporting data in .csv	Exporting data in .csv
Report the extracted data in interactive stereograms	<ul style="list-style-type: none"> • Don't report interactive stereograms. • Necessary export the data collection
<ul style="list-style-type: none"> • The planar facets extracted are easily separated in families and subfamilies sets (interactive) • Spatial orientation similarity (azimuth). 	No interactive data sets
Necessary a good plane exposed to be truly segmented using the tool	Identify planes (with few exposition) and lines using different compass mode tools
Tool advantages: user-friendly (practical, agile and accessible) and free access	Tool advantages: user-friendly (practical, agile and accessible) and free access
<p>Tool disadvantage:</p> <p>Semi-automatic extraction → Slow interpretation process because in this mode a detailed analysis by an expert is necessary to validate the quality and accuracy from the results</p>	Manual extraction → analyst selects the best place to get the measurements.

Comparative table (Facets vs. Compass) according to the results obtained with these two study cases → ARN02 e ARN04

CONCLUSIONS



- The analyst must observe the outcrop wall orientation, as well as the orientation of the structures → These steps are essential to select the best tool to perform the structural analysis of fractures using digital outcrop model
- Positional accuracy and visual quality are crucial for accurate quantitative interpretation of structural features using digital outcrop models, as well as a well-defined data processing routine and careful inspection of the results by an expert.
- The data obtained from this methodological approach will contribute to quantitative analyzes in structural geology based on robust datasets.

ACKNOWLEDGEMENTS

- We would like to thank PETROBRAS (GEOTEC/CENPES/PDEP) (project n° 2017/00057-3) for the financial support to this project.



REFERENCES

Assine, M. L. (2007). Bacia do Araripe. *Boletim de Geociencias Da Petrobras*, 15(2), 371–389.

Dewez, T. J. B., Girardeau-Montaut, D., Allanic, C., & Rohmer, J. (2016). Facets : A cloudcompare plugin to extract geological planes from unstructured 3d point clouds. *International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences - ISPRS Archives*, 41(September), 799–804. <https://doi.org/10.5194/isprsarchives-XLI-B5-799-2016>

Javadnejad, F. (2018). Small Unmanned Aircraft Systems (UAS) for Engineering Inspections and Geospatial Mapping. Thesis. Oregon State University.

Thiele, S. T., Grose, L., Samsu, A., Micklethwaite, S., Vollgger, S. A., & Cruden, A. R. (2017). Rapid, semi-automatic fracture and contact mapping for point clouds, images and geophysical data. *Solid Earth*, 8(6), 1241–1253. <https://doi.org/10.5194/se-8-1241-2017>

Alencar (2016); Brandão (2016); Guimarães (2017); Albuquerque (2017) – Araripe Basin UFPE Mapping