

Atmospheric response to Gulf Stream SST front shifting: impact of horizontal resolution in an ensemble of global climate models

Luca Famooss Paolini^{a,b}, Alessio Bellucci^c, Paolo Ruggieri^{a,d}, Panos Athanasiadis^a, Silvio Gualdi^{a,e}

^a Euro-Mediterranean Centre on Climate Change, Bologna, Italy

^b Ca' Foscari University, Department of Environmental Science, Informatics and Statistics, Venice, Italy

^c Institute of Atmospheric Sciences and Climate - National Research Council (ISAC-CNR), Bologna, Italy

^d University of Bologna, Department of Physics and Astronomy, Bologna, Italy

^e Istituto Nazionale di Geofisica e Vulcanologia, Italy


Contact: luca.famooss@cmcc.it



Università
Ca' Foscari
Venezia



Introduction and objective



Smirnov et al., 2015 - Winter zonally averaged across-front circulation (vectors) and potential temperature (colors) response to a shift in the Oyashio Extension SST front in an atmospheric general circulation model (AGCM). HR – 0.25°; LR – 1°

The present work is a **multi-model analysis** to systematically investigate the **role of horizontal resolution in the atmospheric response** to realistic extratropical SST variability. With this purpose, the atmospheric response to extra-tropical SST anomalies associated with the **inter-annual Gulf Stream SST front (GSF) shifting** during winter has been analyzed.

Data and methods

Table - High Resolution Model Intercomparison Project (HighResMIP) AGCMs. Each model has been forced with the HadISST2 sea ice concentration and SST dataset

Institution	Model	Nominal Resolution (km)	Members
EC-Earth-Consortium	EC-Earth3P	100	3
	EC-Earth3P-HR	50	3
MOHC	HadGEM3-GC31-MM	100	3
	HadGEM3-GC31-HM	50	3
ECMWF	ECMWF-IFS-LR	50	8
	ECMWF-IFS-HR	25	6

“North” phase of the GSF → **Upper tercile** of the GSF latitude
averaged in [50°–68°W]


“South” phase of the GSF → **Lower tercile** of the GSF latitude
averaged in [50°–68°W]

Atmospheric response to GSF shifting



“North” phase minus **“South” phase**


Results – GSF latitude time-series




- Dashed black line is the 10-years running mean

Results – SST anomalies

“North” phase



“South” phase




- **Tripolar structure** extending to the entire North Atlantic in both GSF phases but of opposite sign
- SST anomalies particularly strong close to the GSF winter climatological position



**NAO forcing
&
Intrinsic oceanic processes**

Results – Atmospheric surface response




AGCMs with horizontal resolution = 100 km

- **Negative SLP anomalies** downstream the diabatic heating source, recalling cold air from higher latitude → as expected in «theoretical linear models»
- **Strong surface heat fluxes (SHF)** anomalies close to the GSF winter climatological position

AGCMs with horizontal resolution ≥ 50 km

- **Positive SLP anomalies** downstream the diabatic heating source, recalling warm air from lower latitude
- **Strong SHF anomalies** close to the GSF winter climatological position
- Response comparable to **observations**

Results – Meridional air temperature gradient at 925hPa




AGCMs with horizontal resolution = 100 km

- **Enhancement of meridional air temperature gradient (baroclinicity)** close to the GSF, consistent with SHF anomalies

AGCMs with horizontal resolution ≥ 50 km

- **Enhancement of baroclinicity** close to the GSF, consistent with SHF anomalies
- **Large-scale baroclinicity anomalies** extending downstream and north the GSF (*see next slide*)
- Response comparable to **observations**

Results – Meridional eddy heat fluxes ($v'T'$) at 850hPa




AGCMs with horizontal resolution = 100 km

- **Reduction of meridional eddy heat fluxes (MEHF)** in the western North Atlantic
- Reduction of transient eddy activity also in eastern North Atlantic (*see the supplementary material – Meridional momentum flux at 250hPa*)

AGCMs with horizontal resolution ≥ 50 km

- **Intensification of MEHF** that relaxes the local enhancement of baroclinicity \rightarrow «Atmospheric baroclinic adjustment»
- **MEHF convergence and poleward warm temperature advection** by anomalous mean flow extend the baroclinicity anomalies north and downstream
- Response comparable to **observations**

Results – Zonal wind at 850hPa



AGCMs with horizontal resolution = 100 km

- **Equatorward shifting** of the jet stream
- Increasing in frequency of southern jet position (*see the supplementary material – PDFs of jet stream latitude*)

AGCMs with horizontal resolution ≥ 50 km


- **Poleward shifting** of the jet stream
- Increasing in frequency of northern jet position (*see the supplementary material – PDFs of jet stream latitude*)
- Response comparable to **observations**

Results – Thermodynamic budget

ZONAL TERMS	MERIDIONAL TERMS	VERTICAL TERMS
<div style="border: 1px solid blue; padding: 10px; margin-bottom: 10px;"> <p style="text-align: center;">Mean Thermal Zonal Advection</p> $\bar{u} \frac{\partial \bar{T}}{\partial x} + \frac{\partial}{\partial x} \overline{u' T'}$ </div> <div style="text-align: center;"> <p style="margin-top: 10px;">Zonal Eddy Heat Flux Divergence</p> </div>	<div style="border: 1px solid blue; padding: 10px; margin-bottom: 10px;"> <p style="text-align: center;">Mean Thermal Meridional Advection</p> $\bar{v} \frac{\partial \bar{T}}{\partial y} + \frac{\partial}{\partial y} \overline{v' T'}$ </div> <div style="text-align: center;"> <p style="margin-top: 10px;">Meridional Eddy Heat Flux Divergence</p> </div>	<div style="border: 1px solid blue; padding: 10px; margin-bottom: 10px;"> <p style="text-align: center;">Mean Thermal Vertical Advection</p> $\left(\bar{\omega} \frac{\partial \bar{T}}{\partial p} - \frac{\kappa}{p} \bar{\omega} \bar{T} \right) + \left(\overline{\omega' \frac{\partial T'}{\partial p}} - \frac{\kappa}{p} \overline{\omega' T'} \right)$ </div> <div style="text-align: center;"> <p style="margin-top: 10px;">Vertical Eddy Heat Flux Divergence</p> </div>
$= \bar{Q}$		

- Equation applied to each GSF phase
- Averaged within $\pm 1^\circ\text{N}$ band (band of positive SST anomalies) respect the GSF

Results – Thermodynamic budget




AGCMs with horizontal resolution = 100 km (LR)

- Near-surface diabatic heating anomalies mainly balanced by **mean thermal meridional advection** (cold air from higher latitudes)
- **Mean thermal vertical advection** quite relevant in the interior of the atmosphere (*see next slide*)

AGCMs with horizontal resolution ≥ 50 km (HR)

- Near-surface diabatic heating anomalies balanced by **meridional eddy heat flux divergence**
- **Mean thermal vertical advection** quite relevant in the interior of the atmosphere (*see next slide*)

Results – Omega



AGCMs with horizontal resolution = 100 km

- **Intense upward motion anomalies** in the vicinity of the GSF
- **Equatorward motion** extending throughout the troposphere


AGCMs with horizontal resolution ≥ 50 km

- **Intense upward motion** anomalies in the vicinity of the GSF
- **Weak circulation cell** south of the GSF
- Response comparable to **observations**

Conclusions

- The role of **horizontal resolution** on the atmospheric response to the **interannual GSF shifting** has been investigated in a multi-model multi-member ensemble of atmosphere-only historical simulations
- The atmospheric response to the GSF shifting is **strongly-resolution dependent**
- AGCMs with horizontal resolution = 100 km:
local SHF anomalies → **southward cold air advection** → southward shifting of jet stream
- AGCMs with horizontal resolution ≥ 50 km:
local SHF anomalies → **northward transient eddy heat transport** → large-scale baroclinicity anomalies → northward shifting of jet stream
- AGCMs with horizontal resolution ≥ 50 km are in agreement with observations
- Possible existence of a **positive feedback** between the GSF and NAO:
Positive NAO → poleward GSF shift → positive NAO








Supplementary material – Meridional momentum flux ($v'v'$) at 250hPa



Contours – Winter meridional momentum flux climatology

Shades – Meridional momentum flux anomalies

Black points represent areas statistically significant at the 90% confidence level




AGCMs with horizontal resolution = 100 km

- Reduction of storm track activity in the North Atlantic

AGCMs with horizontal resolution ≥ 50 km

- Poleward shifting of storm track
- Response comparable to **observations**

Supplementary material – PDFs of jet stream latitude



AGCMs with horizontal resolution = 100 km

- PDFs more (less) pronounced towards **southern** (northern) jet position
- Changes in variability of jet position consistent with equatorward jet stream shifting

AGCMs with horizontal resolution ≥ 50 km

- PDFs more (less) pronounced towards **northern** (southern) jet position
- Changes in variability of jet position consistent with equatorward jet stream shifting
- Response comparable to **observations**