


Texture Evolution of Amphiboles - a Case Study from the Mamonia Complex, Cyprus

Amir Topaz, Tzahi Golan, and Yuval Boneh

Ben Gurion University of the Negev, Israel



Previous studies



Soret et al., 2019

Previously suggested deformation mechanisms for amphibole CPO:


Brittle, Cataclastic, Rigid Body Rotation, Grain Growth, Dissolution-Precipitation, Diffusion Creep, Dislocation Creep

Imon et al., Diaz Aspiroz et al., 2006, Pearce et al., 2011, Soret et al., 2019

Unlike many other minerals, CPO in amphibole does not compel Dislocation Creep


Amphibolite samples from the Mamonia complex

P-T: ~600 °C, 6 kbar




<http://www.clima-project.eu/case-studies/nea-paphos/geology/>

Bath of Aphrodite (BOA)




Misorientation map 0-10°

Amphibole Plagioclase




Ayia Varvara (AV)




Crystallographic preferred orientation

Mis-orientation axis



BOA 1



BOA 2



AV




	BOA	AV
Grain Boundaries	Straight	Lobate
Intra-grain orientation spread	✗	✓
Shape orientation affected	✓	✗
Subgrains of (100)[001]	✗	✓
Fractures	✗	✗
Chemical Zoning	✗	✗
<u>Deformation mechanism</u>	Rigid Body Rotation (?) Anisotropic Grain Growth (?)	Dislocation Creep

amirtop@post.bgu.ac.il



Mamonia Complex, Cyprus

- Triassic passive margin
- Comprised of:
 1. volcanic and plutonic rocks (Dhiarizos Group)
 2. sedimentary cover (Ayios Photios Group)
 3. metamorphic outcrops (Ayia Varvara and Loutra tis Aphroditis)
- Juxtaposing the Troodos ophiolite
- Troodos-Mamonia interaction is a serpentinite-decorated strike-slip shear zone
- Several deformation phases were recognized, including subduction-related
- P-T: ~600 °C, 6 kbar







Golan PhD thesis, 2018, modified after Robertson & Xenopontos, 1993 and Swarbrick, 1980.

BOA




AV



Shape parameters:




	BOA		AV	
	XZ	YZ	XZ	YZ
# of grains	13967	4116	5303	1439
long axis diameter (μm)	94	33	63	40
aspect ratio (c:a:b)	3.09:1:0.55		2.44:1:0.59	
shape factor	1.58	1.46	1.56	1.41
paris factor (%)	15.78	28.93	25.44	25.46

$$\text{Shape Factor} = \frac{P}{PE} = \frac{\text{grain perimeter}}{\text{area - equivalent perimeter}}$$





$$\text{Paris Factor} = 2 * \frac{\Delta P}{PE} * 100\%$$

percentile average relative indented surface





CPO and Misorientation

BOA



AV




Crystallographic preferred orientation
*1 point per grain

Misorientation axis (2-10°)




*EBSD data rotated

BOA – two different textures

1.








2.








	plane	# of grains	long axis diameter (μm)	aspect ratio (c:a:b)	shape factor	paris factor (%)
Type 1	XZ	7642	91	3.35:1:0.50	1.62	15.20
	YZ	1933	41		1.44	22.15
Type 2	XZ	6492	88	3.04:1:0.59	1.58	17.15
	YZ	2574	28		1.46	31.75

BOA TYPE 1








BOA TYPE 2




	plane	# of grains	long axis diameter (μm)	aspect ratio (c:a:b)	shape factor	paris factor (%)
Type 1	XZ	7642	91	3.35:1:0.50	1.62	15.20
	YZ	1933	41		1.44	22.15
Type 2	XZ	6492	88	3.04:1:0.59	1.58	17.15
	YZ	2574	28		1.46	31.75


* CPO area-weighted




BOA TYPE 1




BOA TYPE 2




AV




Texture Symmetry



GOS – Grain Orientation Spread




Intra-grain misorientation map



% GOS over threshold (0.035)

GOS map



GOS> threshold map

