



UNIVERSITY OF HELSINKI
FACULTY OF SCIENCE

UCSB



COLORADO SCHOOL OF
MINES



ACADEMY
OF FINLAND

Thermodynamic constraints on assimilation of silicic crust by primitive magmas

Jussi S. Heinonen, University of Helsinki; Frank J. Spera, University of California Santa Barbara; Wendy A. Bohrsen, Colorado School of Mines

Question:

How much assimilation of silicate crust is possible by primitive magmas until they are not “primitive” anymore?

Study method:


Thermodynamic and geochemical modeling using the Magma Chamber Simulator

Models:

Parental melts of different composition (komatiitic, picritic, basaltic), age, and tectonic environment assimilate maximum amounts of average crustal materials.

Answer:

1. Upper (or middle) crust: not much, 20–30 wt.%.
 2. Lower crust: A LOT, up to their weight, i.e. ~100 wt.%.
 3. Basaltic magma (MgO < 12 wt.%, SiO₂ < 52 wt.%, Na₂O + K₂O < 5 wt.%)
 - 4. Picritic magma (MgO = 12–18 wt.%, SiO₂ < 52 wt.%)
 - 5. Komatiitic magma (MgO > 18 wt.%, SiO₂ < 52 wt.%)






The Magma Chamber Simulator (MCS) is a numerical code that quantifies the **thermodynamic** and **geochemical** effects of magma **Recharge**, **Assimilation of partial melts** and/or **Stoped blocks**, and **Fractional Crystallization (RASFC)** on the evolution of a magma body and surrounding wallrock.

To maximize potential for assimilation:

- Wallrock preheated above solidus but below percolation threshold (10 wt.% of melt) temperature
 - LC = 1060 °C @ 800 MPa
 - MC = 880 °C @ 500 MPa
 - UC = 700 °C @ 200 MPa

Assimilation by **stopping** (bulk) vs. wallrock **partial melting** (selective)

- Stopped blocks are heat sinks that substantially increase the crystallinity in the resident magma
- Assimilation of wallrock partial melts energetically favorable (crystallinity in all presented models below 50 wt.%)



Thermodynamic constraints on assimilation of silicic crust by primitive magmas

Jussi S. Heinonen, University of Helsinki; Frank J. Spera, University of California Santa Barbara; Wendy A. Bohrsen, Colorado School of Mines

Parental melts ($\text{SiO}_2 < 52$ wt.%):

Komatiitic/meimechitic ($\text{MgO} > 18$ wt.%)

Etendeka LIP
Karoo LIP (MEI)
Siberian Traps (MEI)
Gorgona Island
Emeishan LIP
Belingwe GB
Abitibi GB
Barberton GB
Comondale GB
Norseman-Wiluna GB

Picritic ($\text{MgO} = 12-18$ wt.%)

Gorgona Island
Grassy Portage Bay GB
NAIP
Hawaii
Karoo LIP
Ontong Java Plateau
Ferrar LIP
Deccan LIP
Siberian Traps
Siberian Traps 2

Basaltic/picrobasaltic ($\text{MgO} = <12$ wt.%)

Vanuatu arc
Early Central American arc
Kurile-Kamchatka arc
Iceland
SWIR MORB
Hawaii tholeiite
Pyrolite model melt
Aleutian arc
Karoo LIP picrobasalt
Ethiopian-Yemeni CFB

Wallrocks

Lower crust (avg)
basaltic andesitic
 $\text{SiO}_2 = 53$ wt.%
at 800 MPa

Middle crust (avg)
dacitic
 $\text{SiO}_2 = 63$ wt.%
at 500 MPa

Upper crust (avg)
dacitic
 $\text{SiO}_2 = 66$ wt.%
at 200 MPa

30 x **3** models = 90 altogether

Rudnick & Gao (2003), Treatise on Geochemistry 3, 1-64.



UNIVERSITY OF HELSINKI
FACULTY OF SCIENCE

UCSB




COLORADO SCHOOL OF
MINES



ACADEMY
OF FINLAND

Thermodynamic constraints on assimilation of lower crust by primitive magmas

Jussi S. Heinonen, *University of Helsinki*; Frank J. Spera, *University of California Santa Barbara*; Wendy A. Bohrsen, *Colorado School of Mines*





UNIVERSITY OF HELSINKI
FACULTY OF SCIENCE

UCSB




COLORADO SCHOOL OF
MINES



ACADEMY
OF FINLAND

Thermodynamic constraints on assimilation of upper crust by primitive magmas

Jussi S. Heinonen, *University of Helsinki*; Frank J. Spera, *University of California Santa Barbara*; Wendy A. Bohrsen, *Colorado School of Mines*





UNIVERSITY OF HELSINKI
FACULTY OF SCIENCE

UCSB




COLORADO SCHOOL OF
MINES



ACADEMY
OF FINLAND

Thermodynamic constraints on assimilation of middle crust by primitive magmas


Jussi S. Heinonen, *University of Helsinki*; Frank J. Spera, *University of California Santa Barbara*; Wendy A. Bohrsen, *Colorado School of Mines*



Thermodynamic constraints on assimilation of silicic crust by primitive magmas

Jussi S. Heinonen, University of Helsinki; Frank J. Spera, University of California Santa Barbara; Wendy A. Bohrsen, Colorado School of Mines

An additional look on the models in SiO₂ vs. MgO space and implications for the generation of siliceous high-Mg basalts (SHMB)



SHMB classification: Pearce & Reagan (2019), Geosphere 15, 1008-1037.



Thermodynamic constraints on assimilation of silicic crust by primitive magmas

Jussi S. Heinonen, *University of Helsinki*; Frank J. Spera, *University of California Santa Barbara*; Wendy A. Bohrson, *Colorado School of Mines*

CONCLUSIONS

- It is difficult for any subalkaline primitive magma to assimilate more than 20–30 wt.% of UC or MC before evolving to compositions with higher SiO₂ than a basaltic magma (52 wt.%).
 - Basaltic magmas cannot assimilate any silicate crustal materials in excess of 20–30 wt.% and remain basaltic
 - SHMBs generated by assimilation of UC/MC by komatiitic magma
- Typical komatiitic and picritic magmas have thermodynamic potential to assimilate considerable amounts (60–100 wt.% and 30–50 wt.%, respectively) of LC and retain a basaltic composition.
 - Generation of basaltic magmas with considerable crustal overprint (e.g., massif-type anorthosites)?
 - Generation of large-scale mineralization fronts related to mafic magmas in lower crust?
 - Crustal growth vs. crustal recycling!
- **Thermodynamic models, as presented here, provide a framework over which additional kinetic and dynamic constraints can be placed to study natural systems in more detail.**



Publicly available for Mac and PC: <https://mcs.geol.ucsb.edu/code>
(or google “Magma Chamber Simulator”)

Thermodynamic constraints on assimilation of silicic crust by primitive magmas

Jussi S. Heinonen, *University of Helsinki*; Frank J. Spera, *University of California Santa Barbara*; Wendy A. Bohrson, *Colorado School of Mines*

MCS presentations in this conference:

SC5.20 **EDI**★

Showcasing the Magma Chamber Simulator▶

Co-organized by GMPV

Convener: Jussi Heinonen  | Co-conveners: Wendy Bohrson , Riikka Fred , Frank Spera , Ville Virtanen 

★ Tue, 27 Apr, 16:00–17:00 (CEST) 

Short course



EGU21-6922 | vPICO presentations | [GMPV8.1/EMRP1.10/NH2.8](#) ★
The Origin and Melt Evolution of Massif-type Anorthosite Parental Magmas: Thermodynamically Controlled Major Element Constraints▶

Riikka Fred, Aku Heinonen, and Jussi S. Heinonen
Mon, 26 Apr, 11:16–11:18



EGU21-2570 | vPICO presentations | [GMPV8.1/EMRP1.10/NH2.8](#) ★
Modelling the production of linear trends in granitoids using the Magma Chamber Simulator: a case study of the Jindabyne Suite from the Lachlan Fold Belt, Australia▶

Kieran Iles and Jussi Heinonen
Mon, 26 Apr, 11:14–11:16



EGU21-7903 | vPICO presentations | [GMPV5.2/ERE5.5](#) ★
Modeling the sulfide saturation in continuously assimilating magmatic systems with the Magma Chamber Simulator▶

Ville Virtanen, Jussi Heinonen, Nicholas Barber, and Ferenc Molnár
Thu, 29 Apr, 15:35–15:37



E-mail: jussi.s.heinonen@helsinki.fi, Twitter: @JussiSHeinonen