

Ocean heat transport as a driver of sea ice extent in CMIP6 models

Jake Aylmer | David Ferreira | Daniel Feltham

Department of Meteorology, University of Reading, UK

✉ j.r.aylmer@pgr.reading.ac.uk

🐦 [@JakeAylmer](https://twitter.com/JakeAylmer)

Summary of key points:

1. Sea ice extent is anticorrelated with Ocean Heat Transport (OHT) ➡
2. **Northern hemisphere:** increased OHT converges mainly near the Atlantic ice edge ➡
3. **Southern hemisphere:** increased OHT converges uniformly under the Antarctic ice pack ➡
4. The different hemisphere behaviours are robust across models ➡
5. Our results suggest ocean biases could contribute substantially to sea ice projection uncertainty ➡

1. Sea ice decreases with OHT

- CMIP6 pre-industrial control simulations
→ correlations (R) based on **25 year means**
- Quantify sea ice cover with zonal-mean **latitude of the ice edge, ϕ_i**
→ we verified this is effectively interchangeable with sea ice extent
- Increased OHT $\Leftrightarrow \phi_i$ shifts poleward
(i.e. sea ice extent decreases)
- Opposite correlation with atmospheric heat transport (AHT)
→ consequence of Bjerknes compensation
→ oceanic component more important on these timescales


Table 1: Bold values are significant correlations at 95% confidence.

Model	Northern Hemisphere		Southern Hemisphere	
	$R(\text{OHT}, \phi_i)$	$R(\text{AHT}, \phi_i)$	$R(\text{OHT}, \phi_i)$	$R(\text{AHT}, \phi_i)$
ACCESS-CM2	+0.86	-0.75	+0.88	-0.74
ACCESS-ESM1-5	+0.94	-0.58	+0.84	-0.83
CAMS-CSM1-0	+0.89	-0.49	+0.94	-0.75
CanESM5	+0.88	-0.71	+0.90	-0.93
CanESM5-CanOE	+0.91	-0.70	+0.92	-0.95
CESM2	+0.73	-0.67	+0.83	-0.54
CESM2-FV2	+0.59	-0.15	+0.65	-0.75
CESM2-WACCM	+0.55	-0.38	+0.90	-0.65
CESM2-WACCM-FV2	+0.82	+0.19	+0.73	-0.66
CNRM-CM6-1-HR	+0.98	-0.83	+0.99	-0.97
CNRM-ESM2-1	+0.98	-0.96	+0.67	-0.71
HadGEM3-GC31-LL	+0.84	-0.74	+0.89	-0.80
HadGEM3-GC31-MM	+0.94	-0.61	+0.96	-0.96
MPI-ESM1-2-HAM	+0.69	-0.56	+0.49	-0.48
MPI-ESM1-2-HR	+0.90	-0.84	+0.71	-0.80
MPI-ESM1-2-LR	+0.77	-0.58	+0.51	-0.52
MRI-ESM2-0	+0.72	-0.32	+0.88	-0.84
NorCPM1	+0.59	-0.46	+0.96	-0.89
UKESM1-0-LL	+0.89	-0.79	+0.88	-0.86

NB: Sign convention

Throughout: OHT/AHT are **poleward** in both hemispheres; and ϕ_i is in **°N** (northern hemisphere) or **°S** (southern hemisphere), so that **each hemisphere may be interpreted symmetrically**

Correlation, $R(x, y)$



1. Sea ice is strongly correlated with OHT

- Models with greater OHT variations (ΔOHT) have larger sea ice variations ($\Delta\phi_i$):




Figure 1

2. What processes underly the Arctic OHT- ϕ_i relationship?

- Difference between maximum and minimum 25 year mean sea ice states

→ HadGEM3-GC31-LL shown here for example

→ Other models behave similarly





Figure 2

3. What processes underly the Antarctic OHT- ϕ_i relationship?

- Difference between maximum and minimum 25 year mean sea ice states

→ HadGEM3-GC31-LL shown here for example

→ Other models behave similarly



4. Robustness across models

A	ACCESS-CM2	F	CESM2	K	CNRM-ESM2-1	P	MPI-ESM1-2-LR
B	ACCESS-ESM1-5	G	CESM2-FV2	L	HadGEM3-GC31-LL	Q	MRI-ESM2-0
C	CAMS-CSM1-0	H	CESM2-WACCM	M	HadGEM3-GC31-MM	R	NorCPM1
D	CanESM5	I	CESM2-WACCM-FV2	N	MPI-ESM-1-2-HAM	S	UKESM1-0-LL
E	CanESM5-CanOE	J	CNRM-CM6-1-HR	O	MPI-ESM1-2-HR		




Figure 4

Averaged over sea ice

- In the Arctic:** increased OHT is mainly lost near the ice edge
 - OHTC under sea ice does not change much (Fig. 4 left)
 - Second effect: increased pan-Arctic AHTC and associated thinning
- In the Antarctic:** increased OHT mainly converges under sea ice and drives basal melt




Figure 5

$\Delta\phi_i$ (sea ice edge)

5. Implications for climate projections




Figure 6: CMIP6 projections of sea ice extent (annual mean) under four emission scenarios (each line: multi-model mean of ensemble means of 15 models; shading indicates one standard deviation). Observations from NSIDC.

- Latest generation models exhibit intermodel spread in future projections (Fig. 6)
 - e.g. size of spread is comparable to differences in emission scenario means

- Our results suggest ocean biases could contribute to sea ice projection uncertainty
 - future work will investigate links between projected rates of sea ice loss and simulated OHT biases (Fig. 7)




Figure 7: OHT in historical simulations of CMIP6 models (same set as in Fig. 6).