

EGU21-3473

Download full paper from the British Columbia Geological Survey website:

http://cmscontent.nrs.gov.bc.ca/geoscience/PublicationCatalogue/Paper/BCGS_P2021-01-08_Rukhlov.pdf

Near-surface mercury vapour haloes in air above ore deposits and faults on Vancouver Island: Insights into buried materials in real-time?



Alexei S. Rukhlov^{1,a}, Luke Ootes¹, Adrian S. Hickin¹, and Nikolay R. Mashyanov²

¹ British Columbia Geological Survey, Ministry of Energy, Mines and Low Carbon Innovation, Victoria, BC, V8W 9N3

² Lumex-marketing LLC, 1 lit. B, Obruchevykh Street, St. Petersburg, 195220, Russia

^a corresponding author: Alexei.Rukhlov@gov.bc.ca

Recommended citation: Rukhlov, A.S., Ootes, L., Hickin, A.S., and Mashyanov, N.R., 2021. Near-surface mercury vapour haloes in air above ore deposits and faults on Vancouver Island: Insights into buried materials in real-time? In: Geological Fieldwork 2020, British Columbia Ministry of Energy, Mines and Low Carbon Innovation, British Columbia Geological Survey Paper 2021-01, pp. 113-143.

Abstract

Volatile geogenic components, such as CO₂, He, Rn, and Hg, form haloes in soil gas and near-surface air directly above mineral deposits. This contrasts with lithochemical, hydrochemical, and biochemical dispersion haloes that can be laterally displaced or obscured by transported overburden. Mercury vapour surveys have been used in geochemical exploration, because Hg occurs in most types of endogenic ore deposits and is highly mobile. In this study, we measured Hg vapour in air 1-50 cm above ground at 15 sites on Vancouver Island. To evaluate the effectiveness of the method across a range of settings, these sites include different types of known mineralized zones, barren rocks, and faults, both buried and exposed. The direct and continuous analysis via a portable RA-915M mercury analyzer reveals Hg vapour concentrations ranging from 0.5 to 54.4 ng·m⁻³. The highest Hg concentration was observed above tailings at the Bentley Au occurrence, possibly due to the amalgamation technique used for fine gold extraction between late 1800s and early 1900s. Prominent Hg vapour haloes mark shear-hosted Cu-Ag-Au sulphides at Mount Skirt (13.4x background Hg), epithermal Au-Ag-Cu at Mount Washington (8.9x background Hg), and sediment-covered polymetallic volcanogenic massive sulphide at the Lara-Coronation occurrence (4.2 to 6.6x background Hg). Basalt-hosted Cu-Ag-Au sulphide zones at the Sunro past producer are marked by weak Hg vapour anomalies relative to local background. Faults, including the Leech River fault, which was active in the Quaternary, are also marked by weak Hg vapour anomalies. The study confirms that, although the Hg level is influenced by weather, the real-time Hg vapour measurement of near-surface air can instantly delineate mineralized zones and fault structures that are buried under overburden 10s of m thick. In contrast to soil gas sampling, this simple and rapid technique can be applied to mineral exploration and geological mapping under overburden above any type of surface, including outcrops, talus, bogs, water bodies, snow, and permafrost.

Keywords: Mercury vapour haloes, near-surface air, real-time analysis, portable RA-915M mercury analyzer, overburden, Leech River fault, Vancouver Island, mineral exploration under cover, polymetallic volcanogenic massive sulphide, epithermal Au-Ag-Cu, shear-hosted sulphide Cu-Ag-Au, Lara-Coronation, Mount Washington Copper, Sunro, Bentley Au, Ralph
