


U-depleted versus U-enriched signatures in complementary crust-mantle sources of volcanic rocks from Asia

Sergei Rasskazov^{1,2},
Irina Chuvashova^{1,2},
Tatiana Yasnygina¹,
Elena Saranina¹

¹Institute of the Earth's Crust SB,
Irkutsk, RF

²Irkutsk State University, Irkutsk, RF



This work is supported by the
RSF grant 18-77-10027


Tian-Shan


Cretaceous-Paleogene
Suluterek Formation

Oligocene-Miocene
Shamsi Series

Jurassic sediments




Minkush












TAS diagram (a), Mg# vs. $\text{CaO}/\text{Al}_2\text{O}_3$ (b), Mg# vs. $10^2 \times \text{K}_2\text{O}/\text{Al}_2\text{O}_3$ (c) and $(\text{Ne}+\text{Lc})-(\text{Hy}+\text{Qz})$ vs. $100 \times \text{An}/(\text{An}+\text{Ab})$

1–6 – groups of K-Pg rocks from Northern Tian Shan: 1 – foidites from Kenkol, 2 – basanites from Uchkuduk, 3 – basanites from Karatayun, Kargon and Kastek, 4 – trachybasalts from Tashkumyr, 5 – basalts and trachybasalts from Suluterek, Bailamtal, and Toraigyr, 6 – basaltic andesites from Tossor, 7 – high-Ti basalts from Tekelik; 8 – K-Pg microbasalt-basanite-phonolite series from the Tuyon basin, data from (Sobel, Arnaud, 2000; Ji et al., 2006; Han et al., 2008).









Th/U



Th/U – Ce/Pb in volcanic
rocks from Southern
Primorye







Comparison of basic pseudotachylytes from the MSF and volcanic rocks from the KSZHT on $(\text{La/Yb})_N - \text{Th/U}$ (a) and $(\text{La/Yb})_N - \text{Nb/U}$ (b) diagrams. Symbols are as in Figure 9. Green arrows indicate extraction of U from layers by oxidized fluids that result in decrease of $(\text{La/Yb})_N$ due to locally increasing degrees of melting. The upper Th/U limit of 20.6 is shown for a source region from Pb isotope data. $(\text{La/Yb})_N$ is calculated using the pyrolite composition after [McDonough, Sun, 1995].

CONCLUSION

Ranges of Th/U ratios of volcanic rocks from the Tunka Valley, Central Baikal Rift System, Kamar-Stanovoi Transtension Zone (KSZHT) in comparison with those from Crust-Mantle Transition sources in other Asian regions

Region, area	Central BRS KSZHT	Sea of Japan Nakhodka v.f.	Tian Shan South Middle-North		Lesser Khingan Udurchukan v.f.
Crust	$2.5 < \text{Th}/\text{U} < 6$	$4 < \text{Th}/\text{U} < 10$	No erupted melts	$0.17 < \text{Th}/\text{U} < 1.8$	No erupted melts
CMT Grt- Grt+ Deeper mantle	$2.5 < \text{Th}/\text{U} < 8$	$5 < \text{Th}/\text{U} < 20$	$2 < \text{Th}/\text{U} < 15.5$	No erupted melts	$4 < \text{Th}/\text{U} < 16$
	$2 < \text{Th}/\text{U} < 10$	No erupted melts	No erupted melts	OIB-like	$8 < \text{Th}/\text{U} < 13$
	$2 < \text{Th}/\text{U} < 6$				OIB-like