

Properties of A Supercritical Quasi-Perpendicular Interplanetary Shock Propagating in Super-Alfvénic Solar Wind: from MHD to Kinetic Scales

Mingzhe Liu^{1,2,3}, Zhongwei Yang², Ying D. Liu^{2,4}, Bertrand Lembège³, Karine Issautier¹, L. B. Wilson III⁵, Siqi Zhao⁶, Vamsee Krishna Jagarlamudi⁷, Xiaowei Zhao²

EGU21-4908

LESIA l'Observatoire de Paris | PSL



1. LESIA, Observatoire de Paris, Université PSL, CNRS, 92195, Meudon, France; mingzhe.liu@obspm.fr;
2. State Key Laboratory of Space Weather, NSSC, CAS, Beijing 100190, China
3. LATMOS/IPSL, UVSQ Paris-Saclay University, Sorbonne University, CNRS, Guyancourt, France
4. University of Chinese Academy of Sciences, Beijing 100049, China
5. NASA Goddard Space Flight Center, Code 672, Greenbelt, Maryland, MD20707, USA
6. Deutsches Elektronen Synchrotron (DESY), Platanenallee 6, D-15738 Zeuthen, Germany
7. National Institute for Astrophysics- Institute for Space Astrophysics and Planetology, 00133 Roma, Italy



1. Abstract

We investigate the properties of an interplanetary shock ($M_A=3.0$, $\theta_{Bn}=80^\circ$) propagating in Super-Alfvénic solar wind observed on September 12th 1999 with *in situ* Wind/MFI and Wind/3DP observations. Key results are obtained concerning the possible energy dissipation mechanisms across the shock and how the shock modifies the ambient solar wind at MHD and kinetic scales: (1) Waves observed in the far upstream of the shock are incompressional and mostly shear Alfvén waves. (2) In the downstream, the shocked solar wind shows both Alfvénic and mirror-mode features due to the coupling between the Alfvén waves and ion mirror-mode waves. (3) Specularly reflected gyrating ions, whistler waves, and ion cyclotron waves are observed around the shock ramp, indicating that the shock may rely on both particle reflection and wave-particle interactions for energy dissipation. (4) Both ion cyclotron and mirror mode instabilities may be excited in the downstream of the shock, since the proton temperature anisotropy touches their thresholds due to the enhanced proton temperature anisotropy. (5) Whistler heat flux instabilities excited around the shock give free energy for the whistler precursors, which help explain the isotropic electron number and energy flux together with the normal betatron acceleration of electrons across the shock. (6) The shock may be somehow connected to the electron foreshock region of the Earth's bow shock, since $B_x > 0$, $B_y < 0$, and the electron flux varies only when the electron pitch angles are less than $PA = 90^\circ$, which should be further investigated.

2. Research background and motivations

2.1 Research background




Fig 1: (Left) Wind Satellite orbit (green) with empirical model bow shock (blue, see Peredo et al., 1995) and magnetopause (red, see Roelof & Sibeck, 1993) plotted for reference. The triangle and diamond respectively represent the start and end of Wind trajectory. The asterisk denotes the position where an IP shock was detected by Wind. (Right) The Wind observations which corresponds to the trajectory shown in the left. From top to bottom, the panels show the magnetic field magnitude, GSE components of the magnetic field, pitch-angle distribution of 340 eV electrons.

❖ IP Shock Information Obtained from SDAT (Vinas & Scudder 1986).

- (1) $M=2.1$ & $M_A=3.0$: Supercritical; $\theta_{Bn}=80$ deg: Quasi-perpendicular;
- (2) $|B|$ increases across the shock: Fast shock; $V_{shn}=532.8$ km/s $> V_p$: Forward shock.

❖ The IP shock was immersed in the quasi-parallel region of the Earth's bow shock.

- (1) Time: 03:57:56 UT on September 12th, 1999
- (2) Location: GSE (33.1, -38.1, -3.9) R_E ; To the Earth's bow shock: ~ 34.0 R_E .

2.2 Motivations

- When IP shocks are Earth-directed, how IP shock interact with the foreshock region of Earth's bow shock? Mutual affection on each other.
- Alfvén waves/fluctuations are usually observed in the foreshock region of Earth's bow shock (eg., Wang et al., 2015 and references therein): Properties of IP shock propagating in the high-Alfvénic solar wind.
- The properties of the bow-shock-reflected electrons and how they are affected by the IP shock.

3. Observations and results

3.1 Overview and MHD properties of the IP shock





Fig 2: Solar wind measurements around the 1999 September 12th shock event from Wind. From top to bottom, the panels show the comparison between magnetic field strength and proton number density, the comparison between X (Y, Z) component of magnetic field and proton bulk velocity in Geocentric Solar Ecliptic (GSE) coordinates, plasma β , the ratio of the perpendicular temperature to the parallel temperature with thresholds of ion cyclotron (red), mirror-mode (blue) and firehose (yellow) instabilities for comparison, the total plasma pressure (proton thermal pressure plus magnetic pressure), and the comparison between cross-helicity and the compressibility, respectively. The red vertical line marks the shock propagating in the ambient solar wind. The region denoted by two blue dashed vertical lines in the downstream of the shock are mirror mode wave structures.



Electron flux from the Sun.

Electron flux reflected by the Earth's bow shock.

Fig 3: Solar wind measurements around the 1999 September 12th shock event from Wind. From top to bottom, the panels show the magnetic field magnitude, proton number density, pitch-angle distribution of 340 eV electrons and electron energy flux at different pitch angles including 14.6°, 34.8°, 56.8°, 78.9°, 101.1°, and 165.4°.

3.2 Kinetic analysis





Fig 5: Wavelet analysis of the magnetic measurements around the 1999 September 12th shock case. From top to bottom, the panels show the magnetic field magnitude, GSE components of the magnetic field, wavelet analysis of magnetic field magnitude $|B|$ and components including B_x , B_y and B_z , respectively.

Time(UT)	$T_{\perp e}/T_{\parallel e}$	$T_{\perp h}/T_{\parallel h}$	$T_{\perp h}/T_{\parallel h}$	$\beta_{\parallel e}$	$n_{he}(cm^{-3})$	n_{he}/n_{se}	n_{he}/n_{se}
03:57:49-03:57:52	0.71	0.65	9.85	1.61	3.61	0.27	0.075
03:57:52-03:57:55	0.72	0.89	6.90	1.67	2.27	0.20	0.087
03:57:55-03:57:58	1.10	1.09	7.56	0.84	4.27	0.27	0.063
03:57:58-03:58:02	1.03	1.10	10.31	1.10	7.35	0.10	0.014
03:58:02-03:58:05	0.98	1.07	10.31	1.23	7.20	0.10	0.015

Table 1: Wind/3DP Electron Parameters from EESA-Low Burst Mode Data.

4. Discussion and perspective

- ✓ Starting from the Earth's bow shock, its electron foreshock region can reach more than 34 R_E .
- ✓ The electrons coming from both the Sun and the Earth's bow shock are pitch-angle scattered across the shock by both the whistler heat flux instabilities and normal betatron acceleration.
- ✓ The IP shock modifies the properties of the waves/fluctuation at the Earth's foreshock region (eg., propagation direction, compression).
- ✓ The evolution of the turbulence across the IP shock needs to be further investigated.

Whistler heat flux instabilities

- (1) Unstable when $T_{\perp h}/T_{\parallel h} > 1.01$;
- (2) Stable when $\beta_{\parallel e} \leq 0.25$.