

Marine heatwaves and cold-spells, and their impact on fisheries in the southern North Sea

Sarah Wakelin¹, Bryony Townhill², Georg Engelhard²,
Jason Holt¹ and Richard Renshaw³

¹National Oceanography Centre, UK

²Centre for Environment, Fisheries and Aquaculture Science, UK & the
International Marine Climate Change Centre

³Met Office, UK

EGU General Assembly April 2021

Background

- Extremes in ocean temperature can harm marine life.
- News headlines focus on increased mortality and impacts on fisheries.
- Temperature also affects physiology, reproduction and growth.
- We study short duration extreme temperature events in the North Sea.
- Key question: do these temperature events impact important fish/shellfish stocks?





Methods: ocean temperature data

- Use near-bed temperatures (NBT) from model reanalysis for 1993 to 2019:
 - High quality (assimilates observations)
 - No gaps in time or space.
- Generate “annual cycle” climatology, and the 10th and 90th percentiles, and smooth (31-day window)¹.

Definitions: periods of longer than 5 days with
 NBT > 90th percentile → **marine heatwaves**
 NBT < 10th percentile → **marine cold-spells**


Example for 2018: off North Norfolk, UK
 cold-spells (blue shading); heatwaves (red shading)

NWSHELF_MULTIYEAR_PHY_004_009		
ATLANTIC- EUROPEAN NORTH WEST SHELF- OCEAN PHYSICS REANALYSIS		
MODEL	● ✕ ● ✕	NWS
T bottomT S SSH 3DUV MLD	ⓘ	
0.067 degree x 0.111 degree (24 depth levels)		
From 1993-01-01 to 2019-12-31		
daily-mean,monthly-mean		
MORE INFO 	ADD TO CART 	WMS Sub-setting



Methods: quantify extreme events

Identify heatwaves and cold-spells per ICES rectangle south of 57°N in the North Sea




ICES (1977) Statistical Rectangle Coding System
<http://ices.dk/sites/pub/CM%20Documents/1977/Gen/ICES%20Statistical%20Rectangle%20System%20Gen0377.pdf>

The **event spread** is the % of ICES rectangles and days with extreme temperatures per time period.

Then define

1. extreme heatwave years: event spread > 7% (●)
1998, 2002, 2003, 2006, 2007, 2014 – 2019
2. extreme cold-spell Q1s: event spread > 15% (●)
1994, 1996, 1997, 2010, 2011, 2013, 2018



Results: impact on fisheries (1)

Significant correlations¹ between heatwaves, cold-spells and detrended fisheries catch data² were found for

species	annual heatwaves*	Q1 cold-spells
annual catches correlations		
sole (<i>Solea solea</i>)	0.60	0.46
sea bass (<i>Dicentrarchus labrax</i>)	0.48	0.47
red mullet (<i>Mullus surmuletus</i>)	-0.47	-0.42
European lobster (<i>Homarus gammarus</i>)	0.49	
Q1 (January to March) catch correlations		
edible crab (<i>Cancer pagurus</i>)		-0.45

No correlations were found with heatwaves or cold-spells and detrended catches of cod (*Gadus morhua*), haddock (*Melanogrammus aeglefinus*), mackerel (*Scomber scombrus*), John Dory (*Zeus faber*) or Norway lobster (*Nephrops norvegicus*).

¹Pearson correlation coefficient r, with *P-value* < 0.05; *with 5 year lag



²Cefas/Defra Fisheries Activity Database (<https://www.gov.uk/guidance/fishing-activity-and-landings-data-collection-and-processing>)

Results: impact on fisheries (2)

Change in catch since previous year: comparing 'normal' years with high event-spread years.

Top figure: cold-spell Q1 (January to March)
Reduction in Q1 catches of edible crab.

Bottom figure: annual heatwaves
Increase in European lobster catches 5 years after heatwave years.



Wakelin et al. *in* von Schuckmann et al.
Copernicus Marine Service Ocean State
Report, Issue 5, *Journal of Operational
Oceanography* (in press)

Discussion and conclusions

- Temperature events might not be sufficiently extreme to affect catches of some species.
- Potential impacts of short term temperature extremes impacting on fish catch data include
 - migration to warmer/colder waters,
 - changes to spawning and growth (eg. impacting on fish catch 5 years later),
 - physiological damage, changing behaviour and increasing vulnerability to being caught,
 - mortality.
- Other factors (eg. changes to fishing effort, trophic or foodweb effects) also impact fish catches.
- To our knowledge, this is the first study to identify a link between extreme temperature events and key fish and shellfish stocks in the North Sea.
 - Increased sole, sea bass and European lobster catches occurred five years after annual heatwave years, while red mullet catches decreased.
 - Catches of sole and sea bass increased in years with widespread January to March cold-spells, while catches of edible crabs and red mullet decreased.



National
Oceanography
Centre

Marine heatwaves and cold-spells, and their impact on fisheries
in the southern North Sea

Sarah Wakelin, Bryony Townhill, Georg Engelhard,
Jason Holt and Richard Renshaw

Reference:

Wakelin et al. *in* von Schuckmann et al. Copernicus Marine
Service Ocean State Report, Issue 5, *Journal of Operational
Oceanography* (in press)