



Climate change in the High Mountain Asia simulated with CMIP6 models

Mickaël Lalande

PhD Student 2019-2022

Supervisors : Martin Ménégoz and Gerhard Krinner


Institut des Géosciences de l'Environnement (IGE, Grenoble, France)

EGU General Assembly 2021 — 27/04/2021

1. Introduction


High Mountain Asia (HMA): Introduction

- The Tibetan Plateau (TP): **world's highest plateau** (average elevation 4000m) → influence on **regional and global climate** (e.g., Kutzbach et al., [1993](#))
- Water supply of over **1.4 billion** living downstream (e.g. Immerzeel et al., [2012](#))
- Climatic regimes:
 - winter **westerly disturbances** (WDs)
 - Indian / East Asian **summer monsoon**
- **Warming** over the HMA and TP (Liu et al., [2000](#); Wang et al., [2008](#)) -> **impacts** on permafrost (Yang et al., [2010](#)), glaciers (Yao et al., [2007](#)), water resources (e.g. Immerzeel et al., [2010](#)), etc.
- **Contrasted trends for precipitation and snow cover** (Kang et al., [2010](#))
- **Lack of observations**: western part and high elevation



Use of GCMs (even if coarse spatial resolution ~50-300km) provides a coherent picture of the large-scale temporal and spatial patterns of key variables at a regional scale !

“Cold bias” over Tibetan Plateau



- Cold biases in models from first AMIP experiments over HMA and TP (Mao and Robock, [1998](#))
- Possible explanations: excess precipitation (Lee & Suh, [2000](#)), snow-ice albedo issues (Su et al., [2013](#)), cold biases in T500 due to smoothed topography (Boos and Hurley, [2013](#)), snow cover parameterization and boundary layer (Chen et al., [2017](#)), lack of high-elevation observation stations in the CRU (Gu et al., [2012](#)), etc.


Our study

1. Biases in CMIP6 for near-surface air temperature, total precipitation and snow cover extent?
2. What are the links between the model biases?
3. Do the model biases impact the trends?
4. Projections over the next century?

2. Data and methods

Data and methods

- 26 CMIP6 GCMs simulations for historical period 1979-2014
- 10 CMIP6 models for the future projections: SSP1-2.6, SSP2-4.5, SSP3-7.0 and SSP5-8.5 (O'Neill et al., [2016](#))
- Observations: CRU (Harris et al., [2014](#)), NOAA CDR (Robinson et al., [2012](#)), APHRODITE (Yatagai et al., [2012](#)) and GPCP (Adler et al., [2016](#))
- Reanalyses: ERA-Interim (Dee et al., [2011](#)) and ERA5 (Hersbach et al., [2020](#))
- Domain: High Mountain of Asia (HMA) including the Tibetan Plateau (TP), with elevation higher than 2500 m.asl (red contour)
- 3 subdomains (> 2500 m.asl): Hindu-Kush Karakoram (HK), Himalaya (HM) and Tibetan Plateau (TP)
- Seasons: winter DJFMA (WDs) and summer JJAS (Asian summer monsoon)





Annual climatologies (1979-2014)

3. Historical bias analysis


tas, snc and pr annual cycles

- stronger biases in **winter** for **tas** ($\sim 2/3^{\circ}\text{C}$) and **snc** ($\sim 20\%$) over HMA
- **large snc spread** -> difficulty to simulate snc in complex topography areas
- **ERA5** bias similar to models -> no assimilation $>1500\text{m}$ (Orsolini et al., [2019](#))
- **pr** obs lower than models -> **snow undercatch** issues by rain gauge (e.g. Jimeno-Saez et al., [2020](#))





Spatial biases and metrics



Bias spatial correlation


Annual spatial correlation of bias over HMA from 1979-2014 climatology

	BCC-CSM2-MR	BCC-ESM1	CAS-ESM2-0	CESM2	CESM2-FV2	CESM2-WACCM	CESM2-WACCM-FV2	CNRM-CM6-1	CNRM-CM6-1-HR	CNRM-ESM2-1	CanESM5	GFDL-CM4	GISS-E2-1-G	GISS-E2-1-H	HadGEM3-GC31-LL	HadGEM3-GC31-MM	IPLS-CM6A-LR	MIROC-ES2L	MIROC6	MPI-ESM1-2-HR	MPI-ESM1-2-LR	MRI-ESM2-0	NorESM2-LM	SAM0-UNICON	TaiESM1	UKESM1-0-LL
tas normalized bias	-0.26	0.14	-0.31	0.06	0.22	0.07	0.22	-0.74	-1	-0.64	-0.43	-0.45	-0.1	-0.18	-0.09	-0.21	-0.87	0.19	0.07	-0.11	-0.02	-0.3	0.25	-0.34	-0.2	-0.1
tas bias / snc bias	-0.51	-0.45	-0.21	-0.02	-0.29	0.01	-0.29	-0.5	-0.39	-0.47	-0.53	-0.4	-0.36	-0.35	-0.28	0.16	-0.62	-0.71	-0.58	0.09	-0.23	-0.16	-0.25	-0.18	-0.09	-0.17
tas bias / pr bias	-0.09	-0.22	-0.08	-0.18	-0.21	-0.19	-0.22	0.02	-0.05	-0.02	0.16	-0.16	-0.11	-0.04	-0.04	-0.07	0.02	-0.07	0.02	-0.37	-0.35	-0.24	-0.26	-0.12	-0.14	-0.02
snc bias / pr bias	0.18	0.48	0.41	-0.22	-0.05	-0.18	-0.04	-0.23	-0.38	-0.23	-0.06	0.04	-0.02	0.03	0.05	-0.04	0.06	0.01	-0.31	-0.12	0.1	-0.22	0.13	0.1	0.01	-0.03
tas bias / elevation	-0.41	-0.04	-0.36	-0.28	-0.09	-0.26	-0.1	-0.56	-0.66	-0.55	-0.32	-0.37	-0.34	-0.43	-0.16	-0.09	-0.63	-0.28	-0.52	-0.3	-0.21	-0.42	-0.05	-0.45	-0.34	-0.12
snc bias / elevation	0.63	0.5	0.5	0.53	0.46	0.51	0.44	0.54	0.67	0.53	0.5	0.45	0.46	0.5	0.47	0.32	0.56	0.41	0.56	0.22	0.24	0.44	0.29	0.48	0.39	0.49
pr bias / elevation	0.18	0.43	0.12	-0.13	0.07	-0.12	0.07	-0.15	-0.31	-0.13	-0.05	-0.08	-0.19	-0.18	0.01	-0.28	-0.06	0.03	-0.05	-0.01	0.15	0.01	-0.01	-0.03	-0.12	0.01

- Significant negative correlations between tas and snc biases
- Less obvious for pr (/!\ APHRODITE underestimate solid precip /!\ -> more negative correlation)
- Correlations between tas/snc biases with elevation -> difficulty representing physical processes at high elevation?

Are trends impacted by overall biases?




Historical trends analysis



- No obvious link between model biases and trends
 - Some strongly biased models have trends close to observations
 - On the contrary, some models with little bias have very different trends
 - Except for snow cover in summer -> very small snow cover
- > All available models are kept for projections

4. Projections


HMA annual projection anomalies (relative to 1995–2014 average)



Projections


- annual median 2081-2100 with respect to 1995-2014 average:
 - tas: 1.9 [1.2 to 2.7] °C (SSP1-2.6) to 6.5 [4.9 to 9.0] °C (SSP5-8.5)
 - relative snc: -9.4 [-16.4 to -5.0] % (SSP1-2.6) to -32.2 [-49.1 to -25.0] % (SSP5-8.5)
 - relative pr: 8.5 [4.8 to 18.2] % (SSP1-2.6) to 24.9 [14.4 to 48.1] % (SSP5-8.5)

- snc and pr models variability underestimated?



HMA versus global

Projections anomalies relative to 1995–2014 average HMA versus GLOBAL



- HMA is warming faster as the rest of the world ?
- 10% faster... compared to NH (without Arctic)
- ~ 4% relative snc loss per 1°C GSAT in winter (linear)
- In summer almost all snc disappear in worst scenario (not linear)
- ~ 6 % relative more pr per 1°C GSAT

Take-home message

Take-home message

- Multimodel analysis with **26 CMIP6 GCMs** over **HMA**
- CMIP6 annual multimodel biases (more pronounced in winter for tas and snc):
 - **cold bias** of **-1.9** [-8.2 to 2.9] $^{\circ}\text{C}$
 - **snc overestimated** 12 [-13 to 43] % (or **52** [-53 to 183] % relative)
 - **pr overestimated** 1.5 [0.3 to 2.9] mm.day⁻¹ (or 143 [31 to 281] % relative) **!\\ obs !**
- No obvious link between biases and trends -> **biased models seems able to reproduce past trends**
- Models **resolution** doesn't seem to improve performances! Additional improvements in **parameterizations** seems essential!
- Other variables might be involved... (cloud cover, aerosols, boundary layer, T500,...)
- Annual projections (2081-2100 with respect to 1995-2014 average with 10 GCMs):
 - median **warming** from **1.9 $^{\circ}\text{C}$** to **6.5 $^{\circ}\text{C}$**
 - relative median **snc decrease** from **-9.4 %** to **-32.2 %**
 - relative median **pr increase** from **8.5 %** to **24.9 %**

Future work

Implementation of a **snow cover parameterization** taking into account the variation of **subgrid topography** in LMDZ

Bibliography

References

Abramowitz, M., & Irene, S. (1964). Handbook of Mathematical Functions, pp. 931-995. Retrieved from
http://people.math.sfu.ca/~cbm/aands/abramowitz_and_stegun.pdf

Adler, Robert; Wang, Jian-Jian; Sapiano, Matthew; Huffman, George; Chiu, Long; Xie, Ping Ping; Ferraro, Ralph; Schneider, Udo; Becker, Andreas; Bolvin, David; Nelkin, Eric; Gu, Guojun; and NOAA CDR Program (2016). Global Precipitation Climatology Project (GPCP) Climate Data Record (CDR), Version 2.3 (Monthly). National Centers for Environmental Information. <https://doi.org/10.7289/N56971M6>

Bookhagen, B., & Burbank, D. W. (2010). Toward a complete Himalayan hydrological budget: Spatiotemporal distribution of snowmelt and rainfall and their impact on river discharge. *Journal of Geophysical Research: Earth Surface*, 115(3), 1–25.
<https://doi.org/10.1029/2009JF001426>

Boos, W. R., & Hurley, J. V. (2013). Thermodynamic bias in the multimodel mean boreal summer monsoon. *Journal of Climate*, 26(7), 2279–2287. <https://doi.org/10.1175/JCLI-D-12-00493.1>

Chen, X., Liu, Y., & Wu, G. (2017). Understanding the surface temperature cold bias in CMIP5 AGCMs over the Tibetan Plateau. *Advances in Atmospheric Sciences*, 34(12), 1447–1460. <https://doi.org/10.1007/s00376-017-6326-9>

Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., ... Vitart, F. (2011). The ERA-Interim reanalysis: configuration and performance of the data assimilation system. *Quarterly Journal of the Royal Meteorological Society*, 137(656), 553–597.
<https://doi.org/10.1002/qj.828>

Deser, C., Terray, L., & Phillips, A. S. (2016). Forced and internal components of winter air temperature trends over North America during the past 50 years: Mechanisms and implications. *Journal of Climate*, 29(6), 2237–2258. <https://doi.org/10.1175/JCLI-D-15-0304.1>

References

- De Wekker, S. F. J., & Kossmann, M. (2015). Convective Boundary Layer Heights Over Mountainous Terrain—A Review of Concepts. *Frontiers in Earth Science*, 3(December), 1–22. <https://doi.org/10.3389/feart.2015.00077>
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., & Taylor, K. E. (2016). Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geoscientific Model Development*, 9(5), 1937–1958. <https://doi.org/10.5194/gmd-9-1937-2016>
- Gu, H., Wang, G., Yu, Z., & Mei, R. (2012). Assessing future climate changes and extreme indicators in east and south Asia using the RegCM4 regional climate model. *Climatic Change*, 114(2), 301–317. <https://doi.org/10.1007/s10584-012-0411-y>
- Harris, I., Jones, P. D., Osborn, T. J., & Lister, D. H. (2014). Updated high-resolution grids of monthly climatic observations - the CRU TS3.10 Dataset. *International Journal of Climatology*, 34(3), 623–642. <https://doi.org/10.1002/joc.3711>
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., ... Thépaut, J. (2020). The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146(730), 1999–2049. <https://doi.org/10.1002/qj.3803>
- Immerzeel, W. W., van Beek, L. P. H., & Bierkens, M. F. P. (2010). Climate Change Will Affect the Asian Water Towers. *Science*, 328(5984), 1382–1385. <https://doi.org/10.1126/science.1183188>
- Immerzeel, W. W., & Bierkens, M. F. P. (2012). Asia's water balance. *Nature Geoscience*, 5(12), 841–842. <https://doi.org/10.1038/ngeo1643>
- Immerzeel, W. W., Wanders, N., Lutz, A. F., Shea, J. M., & Bierkens, M. F. P. (2015). Reconciling high-altitude precipitation in the upper Indus basin with glacier mass balances and runoff. *Hydrology and Earth System Sciences*, 19(11), 4673–4687. <https://doi.org/10.5194/hess-19-4673-2015>

References

- Jimeno-Sáez, P., Pulido-Velazquez, D., Collados-Lara, A.-J., Pardo-Igúzquiza, E., Senent-Aparicio, J., & Baena-Ruiz, L. (2020). A Preliminary Assessment of the “Undercatching” and the Precipitation Pattern in an Alpine Basin. *Water*, 12(4), 1061. <https://doi.org/10.3390/w12041061>
- Kang, S., Xu, Y., You, Q., Flügel, W.-A., Pepin, N., & Yao, T. (2010). Review of climate and cryospheric change in the Tibetan Plateau. *Environmental Research Letters*, 5(1), 015101. <https://doi.org/10.1088/1748-9326/5/1/015101>
- Kutzbach, J. E., Prell, W. L., & Ruddiman, W. F. (1993). Sensitivity of Eurasian Climate to Surface Uplift of the Tibetan Plateau. *The Journal of Geology*, 101(2), 177–190. <https://doi.org/10.1086/648215>
- Lee, D. K., & Suh, M. S. (2000). Ten-year east Asian summer monsoon simulation using a regional climate model (RegCM2). *Journal of Geophysical Research Atmospheres*, 105(D24), 29565–29577. <https://doi.org/10.1029/2000JD900438>
- Li, C., Su, F., Yang, D., Tong, K., Meng, F., & Kan, B. (2018). Spatiotemporal variation of snow cover over the Tibetan Plateau based on MODIS snow product, 2001–2014. *International Journal of Climatology*, 38(2), 708–728. <https://doi.org/10.1002/joc.5204>
- Liu, X., & Chen, B. (2000). Climatic warming in the Tibetan Plateau during recent decades. *International Journal of Climatology*, 20(14), 1729–1742. [https://doi.org/10.1002/1097-0088\(20001130\)20:14<1729::AID-JOC556>3.0.CO;2-Y](https://doi.org/10.1002/1097-0088(20001130)20:14<1729::AID-JOC556>3.0.CO;2-Y)
- Mao, J., & Robock, A. (1998). Surface Air Temperature Simulations by AMIP General Circulation Models: Volcanic and ENSO Signals and Systematic Errors. *Journal of Climate*, 11(7), 1538–1552. [https://doi.org/10.1175/1520-0442\(1998\)011<1538:SATSBA>2.0.CO;2](https://doi.org/10.1175/1520-0442(1998)011<1538:SATSBA>2.0.CO;2)
- O’Neill, B. C., Tebaldi, C., van Vuuren, D. P., Eyring, V., Friedlingstein, P., Hurtt, G., ... Sanderson, B. M. (2016). The Scenario Model Intercomparison Project (ScenarioMIP) for CMIP6. *Geoscientific Model Development*, 9(9), 3461–3482. <https://doi.org/10.5194/gmd-9-3461-2016>

References

- Orsolini, Y., Wegmann, M., Dutra, E., Liu, B., Balsamo, G., Yang, K., ... Arduini, G. (2019). Evaluation of snow depth and snow cover over the Tibetan Plateau in global reanalyses using in situ and satellite remote sensing observations. *The Cryosphere*, 13(8), 2221–2239. <https://doi.org/10.5194/tc-13-2221-2019>
- Palazzi, E., von Hardenberg, J., & Provenzale, A. (2013). Precipitation in the Hindu-Kush Karakoram Himalaya: Observations and future scenarios. *Journal of Geophysical Research: Atmospheres*, 118(1), 85–100. <https://doi.org/10.1029/2012JD018697>
- Robinson, David A.; Estilow, Thomas W.; and NOAA CDR Program (2012): NOAA Climate Data Record (CDR) of Northern Hemisphere (NH) Snow Cover Extent (SCE), Version 1. NOAA National Centers for Environmental Information. <https://doi.org/10.7289/V5N014G9>
- Rangwala, I., Sinsky, E., & Miller, J. R. (2013). Amplified warming projections for high altitude regions of the northern hemisphere mid-latitudes from CMIP5 models. *Environmental Research Letters*, 8(2), 024040. <https://doi.org/10.1088/1748-9326/8/2/024040>
- Salunke, P., Jain, S., & Mishra, S. K. (2019). Performance of the CMIP5 models in the simulation of the Himalaya-Tibetan Plateau monsoon. *Theoretical and Applied Climatology*, 137(1–2), 909–928. <https://doi.org/10.1007/s00704-018-2644-9>
- Serafin, S., Rotach, M. W., Arpagaus, M., Colfescu, I., Cuxart, J., De Wekker, S. F. J., ... Zardi, D. (2020). Multi-scale transport and exchange processes in the atmosphere over mountains. In *Multi-scale transport and exchange processes in the atmosphere over mountains*. <https://doi.org/10.15203/99106-003-1>
- Sharma, E., Molden, D., Rahman, A., Khatiwada, Y. R., Zhang, L., Singh, S. P., ... Wester, P. (2019). Introduction to the Hindu Kush Himalaya Assessment. In *The Hindu Kush Himalaya Assessment* (pp. 1–16). https://doi.org/10.1007/978-3-319-92288-1_1
- Smith, T., & Bookhagen, B. (2018). Changes in seasonal snow water equivalent distribution in High Mountain Asia (1987 to 2009). *Science Advances*, 4(1), e1701550. <https://doi.org/10.1126/sciadv.1701550>

References

- Su, F., Duan, X., Chen, D., Hao, Z., & Cuo, L. (2013). Evaluation of the Global Climate Models in the CMIP5 over the Tibetan Plateau. *Journal of Climate*, 26(10), 3187–3208. <https://doi.org/10.1175/JCLI-D-12-00321.1>
- Usha, K. H., Nair, V. S., & Babu, S. S. (2020). Modeling of aerosol induced snow albedo feedbacks over the Himalayas and its implications on regional climate. *Climate Dynamics*, (0123456789). <https://doi.org/10.1007/s00382-020-05222-5>
- Wang, B., Bao, Q., Hoskins, B., Wu, G., & Liu, Y. (2008). Tibetan Plateau warming and precipitation changes in East Asia. *Geophysical Research Letters*, 35(14), L14702. <https://doi.org/10.1029/2008GL034330>
- Wang, X., Yang, M., Wan, G., Chen, X., & Pang, G. (2013). Qinghai-Xizang (Tibetan) Plateau climate simulation using the regional climate model RegCM3. *Climate Research*, 57(3), 173–186. <https://doi.org/10.3354/cr01167>
- Xu, J., Gao, Y., Chen, D., Xiao, L., & Ou, T. (2017). Evaluation of global climate models for downscaling applications centred over the Tibetan Plateau. *International Journal of Climatology*, 37(2), 657–671. <https://doi.org/10.1002/joc.4731>
- Xue, X., Guo, J., Han, B., Sun, Q., & Liu, L. (2009). The effect of climate warming and permafrost thaw on desertification in the Qinghai-Tibetan Plateau. *Geomorphology*, 108(3–4), 182–190. <https://doi.org/10.1016/j.geomorph.2009.01.004>
- Yang, M., Nelson, F. E., Shiklomanov, N. I., Guo, D., & Wan, G. (2010). Permafrost degradation and its environmental effects on the Tibetan Plateau: A review of recent research. *Earth-Science Reviews*, 103(1–2), 31–44. <https://doi.org/10.1016/j.earscirev.2010.07.002>
- Yao, T., Pu, J., Lu, A., Wang, Y., & Yu, W. (2007). Recent glacial retreat and its impact on hydrological processes on the Tibetan Plateau, China, and surrounding regions. *Arctic, Antarctic, and Alpine Research*, 39(4), 642–650.
[https://doi.org/https://doi.org/10.1657/1523-0430\(07-510\)\[YAO\]2.0.CO;2](https://doi.org/https://doi.org/10.1657/1523-0430(07-510)[YAO]2.0.CO;2)

References



- Yao, T., Thompson, L., Yang, W., Yu, W., Gao, Y., Guo, X., ... Joswiak, D. (2012). Different glacier status with atmospheric circulations in Tibetan Plateau and surroundings. *Nature Climate Change*, 2(9), 663–667. <https://doi.org/10.1038/nclimate1580>
- Yatagai, A., Kamiguchi, K., Arakawa, O., Hamada, A., Yasutomi, N., & Kitoh, A. (2012). APHRODITE: Constructing a Long-Term Daily Gridded Precipitation Dataset for Asia Based on a Dense Network of Rain Gauges. *Bulletin of the American Meteorological Society*, 93(9), 1401–1415. <https://doi.org/10.1175/BAMS-D-11-00122.1>

Supplementary materials

CMIP6 models

CMIP6 institute	CMIP6 model	Resolution (lonxlat)	Grid	Calendar	Member	SSP
BCC	BCC-CSM2-MR	1.1°x1.1°	gn	365_day	r1i1p1f1	
	BCC-ESM1	2.8°x2.8°				
CAS	CAS-ESM2-0	1.4°x1.4°	gn	365_day	r4i1p1f1	
NCAR	CESM2	1.2°x0.9°	gn	noleap	r1i1p1f1	
	CESM2-FV2	2.5°x1.9°				
	CESM2-WACCM	1.2°x0.9°				
	CESM2-WACCM-FV2	2.5°x1.9°				
CNRM-CERFACS	CNRM-CM6-1	1.4°x1.4°	gr	gregorian	r1i1p1f2	
	CNRM-CM6-1-HR	0.5°x0.5°				
	CNRM-ESM2-1	1.4°x1.4°				
CCCma	CanESM5	2.8°x2.8°	gn	365_day	r3i1p2f1	
NOAA-GFDL	GFDL-CM4	1.2°x1.0°	gr1	noleap	r1i1p1f1	
NASA-GISS	GISS-E2-1-G	2.5°x2.0°	gn	365_day	r1i1p1f1	
	GISS-E2-1-H					
MOHC	HadGEM3-GC31-LL	1.9°x1.2°	gn	360_day	r1i1p1f3	
	HadGEM3-GC31-MM	0.8°x0.6°				
IPSL	IPSL-CM6A-LR	2.5°x1.3°	gr	gregorian	r1i1p1f1	
MIROC	MIROC-ES2L	2.8°x2.8°	gn	gregorian	r1i1p1f2	
	MIROC6	1.4°x1.4°			r1i1p1f1	
MPI-M	MPI-ESM1-2-HR	0.9°x0.9°	gn	proleptic_gregorian	r1i1p1f1	
	MPI-ESM1-2-LR	1.9°x1.9°				
MRI	MRI-ESM2-0	1.1°x1.1°	gn	proleptic_gregorian	r1i1p1f1	
NCC	NorESM2-LM	2.5°x1.9°	gn	noleap	r2i1p1f1	
SNU	SAM0-UNICON	1.2°x0.9°	gn	noleap	r1i1p1f1	
AS-RCEC	TaiESM1	1.2°x0.9°	gn	noleap	r1i1p1f1	
MOHC, NIMS-KMA	UKESM1-0-LL	1.9°x1.2°	gn	360_day	r1i1p1f2	

Historical bias analysis



Bias spatial correlation (GPCP)


Annual spatial correlation of bias over HMA from 1979-2014 climatology

	BCC-CSM2-MR	BCC-ESM1	CAS-ESM2-0	CESM2	CESM2-FV2	CESM2-WACCM	CESM2-WACCM-FV2	CNRM-CM6-1	CNRM-CM6-1-HR	CNRM-ESM2-1	CanESM5	GFDL-CM4	GISS-E2-1-G	GISS-E2-1-H	HadGEM3-GC31-LL	HadGEM3-GC31-MM	IPSL-CM6A-LR	MIROC-ES2L	MIROC6	MPI-ESM1-2-HR	MPI-ESM1-2-LR	MRI-ESM2-0	NorESM2-LM	SAM0-UNICON	TaiESM1	UKESM1-0-LL
tas normalized bias	-0.26	0.14	-0.31	0.06	0.22	0.07	0.22	-0.74	-1	-0.64	-0.43	-0.45	-0.1	-0.18	-0.09	-0.21	-0.87	0.19	0.07	-0.11	-0.02	-0.3	0.25	-0.34	-0.2	-0.1
tas bias / snc bias	-0.51	-0.45	-0.21	-0.02	-0.29	0.01	-0.29	-0.5	-0.39	-0.47	-0.53	-0.4	-0.36	-0.35	-0.28	0.16	-0.62	-0.71	-0.58	0.09	-0.23	-0.16	-0.25	-0.18	-0.09	-0.17
tas bias / pr bias	-0.03	-0.33	-0.02	-0.08	-0.2	-0.08	-0.21	0.1	0.02	0.07	0.15	-0.05	-0.07	0.03	0.09	0.07	0.05	-0.12	0.15	-0.24	-0.32	-0.1	-0.25	-0.03	-0.08	0.05
snc bias / pr bias	0.21	0.7	0.45	-0.22	-0.02	-0.18	-0.01	-0.26	-0.36	-0.25	0	-0.05	-0.01	-0.01	0.11	0.09	0.08	0.19	-0.38	-0.09	0.15	-0.23	0.27	0.13	0.02	0.02
tas bias / elevation	-0.41	-0.04	-0.36	-0.28	-0.09	-0.26	-0.1	-0.56	-0.66	-0.55	-0.32	-0.37	-0.34	-0.43	-0.16	-0.09	-0.63	-0.28	-0.52	-0.3	-0.21	-0.42	-0.05	-0.45	-0.34	-0.12
snc bias / elevation	0.63	0.5	0.5	0.53	0.46	0.51	0.44	0.54	0.67	0.53	0.5	0.45	0.46	0.5	0.47	0.32	0.56	0.41	0.56	0.22	0.24	0.44	0.29	0.48	0.39	0.49
pr bias / elevation	0.05	0.37	0.05	-0.27	-0.03	-0.26	-0.04	-0.32	-0.44	-0.3	-0.18	-0.24	-0.28	-0.27	-0.17	-0.49	-0.22	-0.15	-0.2	-0.16	0.05	-0.17	-0.17	-0.15	-0.2	-0.15


Historical trends analysis

Observations and multimodel mean (first realization) seasonal trends (1979-2014)


Observation DJFMA




Multimodel DJFMA







Observation JJAS







Multimodel JJAS



tas trends ($^{\circ}\text{C}/\text{dec}$)



snc trends (%/dec)



pr trends (mm/day/dec)