

Spatiotemporal Clustering of Hydrological Droughts in Peninsular India



DST SERB, Funding ID: SRG/000044



Poulomi Ganguli¹, Bhupinderjeet Singh¹, Aparna Raut^{1,2}

27th April; HS2.4.4: EGU21-87


¹Indian Institute of Technology Kharagpur, India

²Technische Universität Dresden Institut für Hydrologie und Meteorologie, Dresden, Germany




Climate Risks: Droughts in India

Geographic Spread of Droughts in past 30 years



Source: Times of India, April 2016

Drop in water levels in 91 reservoir of the country




Source: The Indian Express; Central Water Commission, April 2016

Research Gaps


- Lack of confidence in drought propagation at tropical rain-fed catchments – little to no study on comprehensive understanding on drought stages – onset, propagation & recovery.
- Most analyses are limited to seasonal to sub-annual scale – higher temporal resolution aid in predictive modelling & forecasting efforts.
- Very few studies investigated “drought-rich” & “drought-poor” locations as well as periods for rain-fed basins in India – region is water-stressed & tends to reach a near day-zero condition (Parvatham & Priyadarshini, 2019).




Delineation of Hydrological Droughts in Peninsular Catchments



Space-Time Clustering of Persistent Droughts




Ahmadi & Moradkhani (2019)





Pattern in Drought Onset

Onset is clustered around Aug-Sept.



Onset is clustered around Oct.


Season of the year



Onset during the beginning of monsoon tends to irregular.


- Onset of hydrological drought is associated with failure or delayed arrival of monsoon (June – September) rainfall
- > 60% sites show mean onset of droughts by the end of two monsoon seasons – August & September.

Detection of Homogeneous Drought Regimes



Regime (Spatial Extent)


- 1 (35%)
- 2 (20%)
- 3 (21%)
- 4 (24%)




Criteria to Select Drought Cluster

- Geospatial locations of gauges
- Mean & Extreme drought attributes
- Catchment-specific attributes – baseflow index, area
- Drought seasonality attributes

Dimension Reduction



Identification of Drought Vulnerable Regions



- **Regime 1:** Frequent and severe droughts coincided with a large recovery period – a partial recovery compounded by a severe episode.
- **Regime 4:** Reverse mirror image of R1 – rare events have low recovery length.

Regime 2 is potentially more resilient to drought – less severe droughts have long recovery length – a few outlying events have smaller recovery length.

Summary: Key Insights

- Variable threshold approach is successfully implemented in tropical rain-fed catchments – temporal evolutions of several drought phases – Onset, growth, persistent & recovery are identified.
- Catchments show a marked seasonality in discharge patterns as well as in onset & termination




- Failure/delayed arrival of monsoon leads to the occurrence of hydrological droughts – Onset is clustered during the end of monsoon.
- A spatial Clustering of drought vulnerable areas are identified through Fuzzy approach 
- Regime-1 in Central peninsular India show severe and frequent droughts compounded by large recovery periods



- Partial recovery compounded by severe & frequent drought may put regional Carbon-budget at a risk, endangering ecosystem productivity & resilience.





References

- Ahmadi, B. & Moradkhani, H. Revisiting hydrological drought propagation and recovery considering water quantity and quality. *Hydrological Processes* **33**, 1492–1505 (2019).
- Parvatam, S. & Priyadarshini, S. On Day Zero, India prepares for a water emergency. *Nature India* <https://www.natureasia.com/en/nindia/article/10.1038/nindia.2019.84> (2019).



Spatiotemporal Clustering of Hydrological Droughts in Peninsular India

Thanks!

- Dr. P. Ganguli, pganguli@agfe.iitkgp.ac.in
- Bhupinderjeet Singh, singhbhupinderjeet096@iitkgp.ac.in
- Aparna Raut, aparnaraut@iitkgp.ac.in

Acknowledgement DST SERB



Poulomi Ganguli¹, Bhupinderjeet Singh¹, Aparna Raut^{1,2}

27th April; HS2.4.4: EGU21-87

¹Indian Institute of Technology Kharagpur, India

²Technische Universität Dresden Institut für Hydrologie und Meteorologie, Dresden, Germany

