




High summer precipitation reduces soil methane sink capacity and alters decomposition processes in a mature temperate forest

Katy Faulkner, Simon Oakley, Sally Hilton, Kelly Mason, Sami Ullah, Niall McNamara and Gary Bending

Background



- Temperate forests store ~ 120 Pg carbon (Pan et al., 2011)
- Act as sinks or sources for major greenhouse gases including CO₂, CH₄ and N₂O thus play an important role in global carbon and nitrogen budgets (Friedlingstein et al., 2021; Saunois et al., 2020; Tian et al., 2020)
- Shifting precipitation patterns and increased frequency, duration and magnitude of high rainfall events (Hajek and Knapp, 2022)
- Soil moisture is a key driver of microbial processes (Lui et al., 2009)



Methodology

Research Objectives:


1. Determine the resilience of forest soil **microbial structure** and **function** to **altered rainfall**
2. Determine the **direct** and **indirect** contributions of **soil** and **microbial properties** on soil CO_2 , CH_4 and N_2O fluxes

- Oak (*Quercus robur*) dominated **mature temperate forest** in Staffordshire, UK
- **30% increase** in volumetric soil moisture content for 8-week period
- **Molecular and functional analyses**

(a) *in situ* field experiment




How will high rainfall events impact forest soil microbial function?




- Significant **reduction** in **methane** sink capacity (21 – 67%) during the **high rainfall** treatment period with **limited recovery** 1- and 4-weeks post-treatment
- Significant **reduction** in soil **respiration** (2 - 23%) during the high rainfall period with **no recovery** 1- and 4-weeks post-treatment

How will high rainfall events impact forest soil microbial function?

Hydrolytic





Oxidative





- Reduction in hydrolytic enzyme activity after initiation of simulated high rainfall but fast recovery
- Strongest responses in *N*-acetyl β -glucosaminidase (NAG; chitinase) and peroxidase (perox; lignin degradation)

How does soil moisture drive forest soil CO₂ flux?



- Soil **moisture** indirectly alters the soil CO₂ flux via **increased peroxidase activity** and **reduced microbial biomass N**
- Significant **reduction** in CO₂ flux with increased soil moisture during experiment

How does soil moisture drive forest soil CH₄ flux?



- Soil moisture both **directly** and **indirectly** affects the soil CH₄ flux via methanotroph abundance
- Significant **increase** in CH₄ flux with increased soil moisture during experiment

Summary and acknowledgments




- CO_2 and CH_4 fluxes significantly altered from simulated high rainfall period and do not recover after 4-weeks
- **Soil moisture** primary driver of CH_4 flux which is **highly responsive** to high rainfall treatment but methanogenic and methanotrophic community abundance are poor predictors of net CH_4 flux
- **Soil moisture** indirectly reduces CO_2 flux via altered soil enzyme activities and microbial biomass



Prof. Gary Bending
Dr. Sally Hilton
Dr. Ryan Mushinski

Prof. Yin Chen
Dr. Kevin Purdy
Jessica Webster



Prof. Niall McNamara
Simon Oakley
Kelly Mason

UNIVERSITY OF
BIRMINGHAM



Prof. Sami Ullah
BIFoR Technical team
Prof. Rob MacKenzie

@katyjfaulkner 
katy.faulkner@warwick.ac.uk

