

# A global analysis of water storage variations from remotely sensed soil moisture and daily satellite gravimetry


**Daniel Blank<sup>1</sup>, Annette Eicker<sup>1</sup>, Laura Jensen<sup>1</sup>, and Andreas Güntner<sup>2</sup>**

<sup>1</sup> HafenCity University, Geodesy and Geoinformatics, Hamburg, Germany


<sup>2</sup> Helmholtz-Centre Potsdam - GFZ German Research Centre for Geosciences, Potsdam, Germany

**EGU 2022**

**May 26<sup>th</sup>, 2022**





Satellite observations of surface soil moisture:







ITSG-Grace2018 vs. SMOS L3 SSM



Area-weighted average = 0,18  
Maximum = 0,90





ITSG-Grace2018 vs. SMAP L3 SSM



Area-weighted average = 0,26  
Maximum = 0,95





## ITSG-Grace2018 vs. ESA CCI SSM



Area-weighted average = 0,26  
Maximum = 0,90






ITSG-Grace2018 vs. SMAP L4 SSM



Area-weighted average = 0,52  
Maximum = 0,94



ITSG-Grace2018 vs. SMOS L4 RZSM





- SSM SMOS L3
- SSM SMAP L3
- SSM ESA CCI
- SSM SMAP L4
- RZSM SMOS L4

Area-weighted average = 0,26  
Maximum = 0,98




ITSG-Grace2018 vs. SMAP L4 RZSM




- SSM SMOS L3
- SSM SMAP L3
- SSM ESA CCI
- SSM SMAP L4
- RZSM SMOS L4
- RZSM SMAP L4

Area-weighted average = 0,61  
Maximum = 0,98






















































ITSG-Grace2018 vs. SMAP L4 RZSM






- Global comparison of TWS and soil moisture products
- Calculation of correlations and time-shifts on a global scale
- Signal decomposition into sub-monthly signals



- High correlations between SSM products and TWS. Even stronger correlations with RZSM products.
- TWS is shifted by 1-2 months relative to the SSM and 0-1 month relative to RZSM.
- On sub-monthly time scales, there is agreement between SM and TWS in particular during rainfall events/periods.




**Results:** Blank et al. 2022 (submitted)

*Blank, D., Eicker, A., Jensen, L., Güntner, A., 2022. A global analysis of water storage variations from remotely sensed soil moisture and daily satellite gravimetry*

**Questions?** [daniel.blank@hcu-hamburg.de](mailto:daniel.blank@hcu-hamburg.de)





## Supplementary Slides







# Correlation total signal

Correlation between TWS (ITSG-Grace2018) and satellite soil moisture products







Time-Shift between TWS (ITSG-Grace2018) and satellite soil moisture products
















### ITSG-Grace2018 vs. SMOS L3 SSM



### ITSG-Grace2018 vs. SMAP L4 SSM



### ITSG-Grace2018 vs. SMOS L4 RZSM



### ITSG-Grace2018 vs. ESA CCI SSM

