

Using K-Means Clustering to Compare Adjoint Waveform Tomography Models of California and Nevada

Abstract:




Claire Doody¹, Arthur Rodgers², Michael Afanasiev³, Christian Boehm³, Lion Krischer³, Andrea Chiang², Nathan Simmons²

¹University of California, Berkeley, ²Lawrence Livermore National Laboratory, ³Mondaic Ltd.

This work was supported by LLNL Laboratory Directed Research and Development project 20-ERD-008. This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344. LLNL-PRES-835369


Full waveform inversion (FWI) methods require a starting model, but the effect the starting model has on the final model has not been studied extensively.



Tromp, 2020


We begin with 3 different starting models, each computed on different scales and with different methodologies.

SPIRaL (Simmons et al., 2021)




- Global model
- P/S travel times, Rayleigh wave group velocities, vertical transverse anisotropy

CSEM_NA (Krischer et al., 2018)




- Regional model of North America and Europe
- Adjoint waveform tomography

WUS256 (Rodgers et al., in review)





- Regional model of the Western US
- Adjoint waveform tomography

We use a dataset of 103 events recorded at over 1300 stations.




- Magnitude Range (M_w): 4.5-6.5
- Date Range: Jan 1 2000- Oct 31 2020
- Focal Mechanisms taken from GCMT


We use a dataset of 103 events recorded at over 1300 stations.



Model results show broad similarities in structure, but have differences that can be related to the starting model.





Model results show broad similarities in structure, but have differences that can be related to the starting model.



K-Means clustering analysis confirms the broad structural agreement of all 3 models.

Clustering Analysis of the Upper Mantle on All Models




Clusters of velocity profiles in the crust (0km-30km)

2 Clusters


K-Means clustering analysis confirms the broad structural agreement of all 3 models.

Clustering Analysis of the Upper Mantle on All Models




K-Means clustering analysis confirms the broad structural agreement of all 3 models.


Clustering Analysis of the Upper Mantle on All Models




K-Means clustering analysis confirms the broad structural agreement of all 3 models.




K-Means clustering analysis confirms the broad structural agreement of all 3 models.




K-Means clustering analysis confirms the broad structural agreement of all 3 models.




Waveforms for all three models also show similar levels of fit to the observed data.



Waveforms for all three models also show similar levels of fit to the observed data.



Waveforms for all three models also show similar levels of fit to the observed data.



Questions/Comments:
claired@berkeley.edu

Observed	CANV_SPiRaL	CANV_256	CANV_CSEM
----------	-------------	----------	-----------