
Soil microbial co-occurrence networks and functioning along an aridity gradient in Atlantic coastal dunes

María José Fernández-Alonso, Alexandra Rodríguez,
Raúl Ochoa-Hueso, Fernando T. Maestre, Jorge Durán

1

Introduction Background

• Atlantic coastal dunes


- dynamic and changing ecosystems
- highly sensitive to climate change
- will cross aridity thresholds

→ may endanger ecosystem services ???




Coastal dune site in
Santo André
AI = 0,43

P/PET change (%)



Cherlet et al., 2018 World
Atlas of Desertification



1

Introduction Background

• Atlantic coastal dunes


- dynamic and changing ecosystems
- highly sensitive to climate change
- will cross aridity thresholds

→ may endanger ecosystem services ???




Coastal dune site in
Santo André
AI = 0,43

P/PET change (%)



Cherlet et al., 2018 World
Atlas of Desertification



Biotic cover loss

Reduced nutrient availability

Drastic changes in the composition and structural organization of soil microbes

1

Introduction Aims

- To understand the integrative effects of:

(i) **increasing aridity** & (ii) **biotic cover**
(regional scale) (local scale)

on soil biotic and abiotic attributes and processes in coastal dune ecosystems.

- Microbial abundance and diversity indexes
- Coupling of soil microbial networks

- Water holding capacity
- pH
- Soil organic
- Total nitrogen
- Total phosphorus

- Nitrogen mineralization
- Potential fluxes of greenhouse gases

- High-throughput sequencing (16S, ITS) & qPCR

- Soil physicochemical properties

- Potential soil functions



San Xurxo
AI = 0,92



La Antilla
AI = 0,29

1

Introduction Aims

- To understand the integrative effects of:

(i) **increasing aridity** & (ii) **biotic cover**
(regional scale) (local scale)

on soil biotic and abiotic attributes and processes in coastal dune ecosystems.

- | | |
|---|---|
| <ul style="list-style-type: none"> • Microbial abundance and diversity indexes • Coupling of soil microbial networks • Water holding capacity • pH • Soil organic • Total nitrogen • Total phosphorus • Nitrogen mineralization • Potential fluxes of greenhouse gases | <ul style="list-style-type: none"> • High-throughput sequencing (16S, ITS) & qPCR • Soil physicochemical properties • Potential soil functions |
|---|---|



San Xurxo
AI = 0,92



La Antilla
AI = 0,29

2



Materials and Methods Experimental design

- 24 dune systems distributed along a **natural aridity gradient** (~1500 km) in the Iberian Peninsula (AI, Aridity Index).

$$AI = \frac{\sum_{i=1}^{30} \left(\frac{P_i}{PET_i} \right)}{30}$$

- We carry out a **stratified sampling design** under *microsites*:

Vascular plants Cryptogams Bare soil



2

Materials and Methods Experimental design

- 24 dune systems distributed along a **natural aridity gradient** (~1500 km) in the Iberian Peninsula (AI, Aridity Index).

$$AI = \frac{\sum_{i=1}^{30} \left(\frac{P_i}{PET_i} \right)}{30}$$

- We carry out a **stratified sampling design** under *microsites*:


Vascular plants Cryptogams Bare soil

- We classified dune systems into **3 bioclimatic areas**:

● Humid

■ Transition sites

▲ Drylands



2. Ortigueira



4. San Xurso



12. Dunas São Jacinto



13. Leirosa





16. Santo André



24. Genoveses

3

Results and discussion Climate effects on the biotic cover



Aridity decreased total vegetation cover and favoured the formation of shrub-dominated patches.

Humid

Transition sites

Drylands



2. Ortigueira



4. San Xurso



12. Dunas São Jacinto



13. Leirosa




16. Santo André




24. Genoveses

3

Results and discussion Climate effects on the biotic cover



Humid



Transition sites

Aridity decreased total vegetation cover and favoured the formation of shrub-dominated patches.

Drylands



2. Ortigueira



4. San Xurso



12. Dunas São Jacinto



13. Leirosa




16. Santo André




24. Genoveses

3

Results and discussion Climate effects on the biotic cover



Humid



Transition sites

Aridity decreased total vegetation cover and favoured the formation of shrub-dominated patches.

Drylands



2. Ortigueira



4. San Xurso



12. Dunas São Jacinto



13. Leirosa




16. Santo André



24. Genoveses

3 Results and discussion Climate effects on soil attributes and functioning



Vegetation shifts were accompanied by declines in water holding capacity, organic matter, nutrient contents and N mineralization.

Humid

Transition sites

Drylands



2. Ortigueira



4. San Xurso



12. Dunas São Jacinto



13. Leirosa




16. Santo André



24. Genoveses

3 Results and discussion Climate effects on soil attributes and functioning



Vegetation shifts were accompanied by declines in water holding capacity, organic matter, nutrient contents and N mineralization.

Humid

Transition sites

Drylands



2. Ortigueira



4. San Xurso



12. Dunas São Jacinto



13. Leirosa




16. Santo André



24. Genoveses

3 Results and discussion Climate effects on soil attributes and functioning



Vegetation shifts were accompanied by declines in water holding capacity, organic matter, nutrient contents and N mineralization.

Humid

Transition sites

Drylands



2. Ortigueira



4. San Xurso



12. Dunas São Jacinto



13. Leirosa




16. Santo André



24. Genoveses

3 Results and discussion Climate effects on soil attributes and functioning



Vegetation shifts were accompanied by declines in water holding capacity, organic matter, nutrient contents and N mineralization.

Humid

Transition sites

Drylands



2. Ortigueira



4. San Xurso



12. Dunas São Jacinto



13. Leirosa




16. Santo André










24. Genoveses

3 Results and discussion Climate and microsite effects on soil microbial communities



Soil bacterial abundance was more sensitive to changes in aridity than fungal abundance.



Humid

Transition sites

Drylands



2. Ortigueira

4. San Xurso


12. Dunas São Jacinto

13. Leirosa




16. Santo André

24. Genoveses




3 Results and discussion Climate and microsite effects on soil microbial communities



Soil bacterial abundance was more sensitive to changes in aridity than fungal abundance.



Plant
Biocrust
Bare



Humid

Transition sites

Drylands



2. Ortigueira

4. San Xurso


12. Dunas São Jacinto

13. Leirosa

16. Santo André








24. Genoveses

3 Results and discussion Climate and microsite effects on soil microbial communities



Soil bacterial abundance was more sensitive to changes in aridity than fungal abundance.

Soil fungal communities were more sensitive to the type of microsite present.



Humid

Transition sites

Drylands



2. Ortigueira

4. San Xurso


12. Dunas São Jacinto

13. Leirosa

16. Santo André




24. Genoveses

3 Results and discussion Climate and microsite effects on soil microbial communities






Soil bacterial abundance was more sensitive to changes in aridity than fungal abundance.

Soil fungal communities were more sensitive to the type of microsite present.



Plant
Biocrust
Bare



Humid

Transition sites

Drylands



2. Ortigueira



4. San Xurso



12. Dunas São Jacinto



13. Leirosa




16. Santo André




24. Genoveses

3 Results and discussion Climate and microsite effects on soil microbial communities



Soil bacterial abundance was more sensitive to changes in aridity than fungal abundance.


Soil fungal communities were more sensitive to the type of microsite present.



Humid

Transition sites

Drylands



2. Ortigueira



4. San Xurso



12. Dunas São Jacinto



13. Leirosa





16. Santo André



24. Genoveses


3

Results and discussion Climate effects on soil microbial networks




3

Results and discussion Climate effects on soil microbial networks




Window size =
5 dune sites




3

Results and discussion Climate effects on soil microbial networks



Window size =
5 dune sites



Statistical differences

- p-value < 0.05
- n.s.

Null model


- 95% confidence interval

Linear regressions


- $-0.17 \leq \text{Aridity} \leq 0.44$
- $0.50 \leq \text{Aridity} \leq 0.71$
- $-0.17 \leq \text{Aridity} \leq 0.71$

3

Results and discussion Climate effects on soil microbial networks




Increases in aridity drove a strong increase in the coupling of the soil microbial network until a specific threshold beyond which it remained constant.




3

Results and discussion Climate effects on soil microbial networks



Increases in aridity drove a strong increase in the coupling of the soil microbial network until a specific threshold beyond which it remained constant.



Results and discussion

Microsites as modulators of soil microbial networks

Microsites strongly drove the interactions between soil bacterial groups, but much less so for fungal groups.




Plant



Biocrust



Bare soil



Results and discussion

Microsites as modulators of soil microbial networks

Microsites strongly drove the interactions between soil bacterial groups, but much less so for fungal groups.




Plant



Biocrust



Bare soil



Highlights

- Aridity decreased total vegetation cover and favoured the formation of shrub-dominated patches.
- Aridity negatively impacted soil bacterial abundance, microbial diversity, nutrient contents and potential N mineralization.
- Aridity caused a strong increase in the coupling of the soil microbial networks until a specific threshold ($AI=0,5-0,6$) beyond which it remained constant.
- Soil bacterial communities were more sensitive to changes in aridity than fungal communities.

Acknowledgements:





María José Fernández-Alonso
mjfa@uc.pt

Thanks for your attention!

Acknowledgements:



Next steps

- Incorporation of spatial and environmental information to discern the complex ecological mechanisms driving the microbial community assembly
- Explore for ecological clusters associated to microhabitats: identification of taxa that tend to co-occur (i.e. 'generalists') from those that exclusively occurred in one of the microhabitats (i.e. 'specialists')
- Identify keystone taxa (e.g. aridity sensitive vs. aridity tolerant).

Acknowledgements:

