



L. Macotela^{* (1)}, J. Manninen⁽²⁾, M. Fullekrug⁽¹⁾, I. Kolmasova^(3,4), and O. Santolik^(3,4)

VLF kHz



- Sferics obscure VLF waves



Affiliations

- (1) Department of Electronic & Electrical Eng., U of Bath, UK
- (2) Sodankylä Geophysical Observatory, U of Oulu, Finland
- (3) Institute of Atmospheric Physics, The Czech Academy of Sciences, Czechia
- (4) Faculty of Mathematics and Physics, Charles University, Czechia

Seasonal variability of VLF banded structured events observed in the 5–39 kHz frequency range in Finland

Acknowledgment This project has received funding from the European Union's Horizon 2020 research and under the Marie innovation programme Sklodowska-Curie grant agreement No 101019319

4	* ¹⁰⁴ Waveform · F	K.
2	×10 ⁴	vv
		. اد
В	 Environmentality 	W
-2	×10 ⁴	
-4	×10 ⁴	
1005	0.725	0 Tim



*elmc23@bath.ac.uk

When analysing 1-hr field spectrograms, in the freq. range, several strip elements varying in time and freq. are at times observed. We these events structures

References

Macotela et al., 2019. *Geoph. Res. Let.*, **46**, 24. Titova et al., 2007. Geoph. Res. Let., 34, L02112.