Decadal- to centennial-scale climate changes over the last 2000 yr recorded from varved sediments of Lake Kusai, northern Qinghai-Tibetan Plateau

Weihan Jiaa, Xingqi Liua*

^aCollege of Resource Environment and Tourism, Capital Normal University, Beijing 100048, China *Corresponding author e-mail address: xqliu@cnu.edu.cn (X. Liu).

(RECEIVED June 26, 2018; ACCEPTED March 20, 2019)

Abstract

High-resolution climate archives of the Late Holocene are essential in the study of paleoclimatic dynamics and for understanding the importance of natural and anthropogenic influences on past and future climate changes. Here, we present well-dated Xray fluorescence scanning records retrieved from a varved sediment core from Lake Kusai. These records show the decadal- to centennial-scale paleoclimatic variability of the northern Qinghai-Tibetan Plateau over the last 2000 yr. Ca is mainly related to the precipitation of authigenic carbonates and is a proxy for temperature changes. The Ca record of Lake Kusai is well-correlated with the variations and periodicities of solar activity. Therefore, solar output can be suggested as being the predominant forcing mechanism of decadal- to centennial-scale temperature fluctuations over the last 2000 yr. The evolution of effective moisture was inferred from the log-ratios of Rb/Sr, which demonstrated synchronous changes with the typical Indian summer monsoon record from Dongge Cave. These results indicate that the decadal- to centennial-scale effective moisture evolution of the northern Qinghai-Tibetan Plateau is mainly influenced by the Indian summer monsoon. Additionally, we have not found the evident periodicities of solar activity in our effective moisture record over the last 2000 yr.

Keywords: Lake Kusai; 2000 yr; Decadal to centennial timescales; XRF scanning; Solar activity; Indian summer monsoon; Northern Qinghai-Tibetan Plateau

INTRODUCTION

The last 2000 yr is a crucial period for the study of past global changes (PAGES, 2009; Yan et al., 2015). High-resolution paleoclimatic records for the last 2000 yr are particularly needed to evaluate the increasing impact of human activity and to help us predict the possible evolution of future climate. Accordingly, numerous studies based on stalagmites (Zhang et al., 2008; Tan et al., 2010), lake sediments (Liu, X., et al., 2014a; Li et al., 2017), ice cores (Thompson et al., 2000), tree rings (Shao et al., 2005; Grießinger et al., 2017), and historical documents (Tan et al., 2008) have been performed to shed light on the paleoclimatic changes in China over the last 2000 yr.

The Qinghai-Tibetan Plateau (QTP) is thought to play a key role in the evolution of Asian atmospheric circulation

(An et al., 2001). The paleoclimatic records of the QTP have been well-studied based on lacustrine sediments in different regions (Gasse et al., 1991; Shen et al., 2005; Herzschuh, 2006; Zhu et al., 2008; Herzschuh et al., 2014). These records, however, are mostly concentrated on millennial to orbital timescales, and little is currently known about the paleoclimatic changes of the northern QTP at the decadal to centennial timescales.

Records from Lake Kusai show that paleoclimatic changes at decadal to centennial timescales are mainly influenced by the Asian monsoon and have a close relationship with solar activity (Liu, X., et al., 2009). The chronology obtained from bulk organic carbon, however, is relatively low resolution and is affected by a 3400-yr reservoir effect. To better understand decadal- to centennial-scale paleoclimatic changes and their forcing mechanisms in the northern QTP over the last 2000 yr, we present high-resolution X-ray fluorescence (XRF) scanning records retrieved from a varved sediment core from Lake Kusai.

Cite this article: Jia, W., Liu, X. 2019. Decadal- to centennial-scale climate changes over the last 2000 yr recorded from varved sediments of Lake Kusai, northern Qinghai-Tibetan Plateau. *Quaternary Research* 92, 340–351. https://doi.org/10.1017/qua.2019.19

STUDY REGION

Lake Kusai (approximately 35°37'-35°50'N and 92°35'-93° 05'E, 4475 m above sea level [asl]), a saline inland lake with a present-day salinity of 28.54 g/L, is situated in the Hoh Xil region of the northern QTP, northwest China (Fig. 1A). The area of the lake is 254.4 km², with a catchment of approximately 3700 km². The lake is fed mainly by the Kusai River, which originates to the west of the lake on the Daxue Mountain (5863 m asl). The Kusai River flows into the southwestern part of the lake, which is rather shallow and has a mean water depth of approximately 10 m. The lake is much deeper in the northwest, however, where the depth is approximately 50 m (Li, 1996; Wang and Dou, 1998), and the sedimentary environment is stable and experiences less disturbances than that in the southwest. Modern meteorological records indicate a mean annual temperature of the lake area of approximately -5.4°C, a mean annual precipitation of approximately 275 mm, and a potential annual evaporation of approximately 1300 mm, resulting in a strongly arid climate.

The climate of the northern QTP is affected by both Asian monsoonal systems and the mid-latitude westerly circulation (An et al., 2012; Yao et al., 2013). In the summertime, the lake basin is more humid with atmospheric precipitation and glacial meltwater, both of which can increase the runoff of the Kusai River and enhance the erosional degree of the catchment. In contrast, the region is characterized by a dry and cold continental climate in the winter. The Hoh Xil region experiences the highest frequency of dust storms in the Qinghai Province between February and April (Fang et al., 1997).

MATERIALS AND METHODS

We drilled a 1.52-m-long sediment core (KSD-1) in the northwest of Lake Kusai at a water depth of 49.8 m using the UWITEC platform of Austria (Fig. 1B; http://www.uwitec.at). A short core (17 cm) was also collected parallel to the long core using a gravity corer. All sediment cores were stored in PVC tubes in the field. The short core was sampled at 0.5-cm intervals, and its slices were radiometrically dated by measuring the activity of ²¹⁰Pb and ¹³⁷Cs as a function of depth. The sedimentation rates and chronologies were calculated using the ²¹⁰Pb data and the constant initial concentration (CIC) and the constant rate of supply (CRS) models (Appleby and Oldfieldz, 1983; Oldfield and Appleby, 1984; Appleby, 2001).

Due to the low pollen concentrations and the absence of terrestrial plant macrofossils in the cores, the ¹⁴C dating of total organic carbon samples was performed using accelerator mass spectrometry (AMS). Ten samples from the long sediment core were processed and dated at the National Isotope Centre, Institute of Geological and Nuclear Sciences, New Zealand (Table 1).

The sediment cores of Lake Kusai are rhythmically laminated. The long core was split along the central axis. Undisturbed sediment slabs $(7 \times 1.5 \times 1 \text{ cm})$ were collected continuously down core with 2 cm of overlap between two consecutive slabs to ensure that all couplets of laminations were represented. All sediment slabs were placed in aluminium boxes and frozen quickly by liquid nitrogen in approximately 15 minutes. The slabs were subsequently freeze-dried for 48 hours to remove the residual moisture and then embedded in low-viscosity epoxy resin under low vacuum.



Figure 1. (color online) (A) Map showing the climatic system of China including the Indian Summer Monsoon (ISM), the East Asian Summer Monsoon (EASM), and the Westerlies. The locations of Lake Kusai and other paleoclimate sites mentioned in this paper, including the Guliya ice core (Yao et al., 1996), Dunde ice core (Thompson et al., 2006), Ngamring Tso (Conroy et al., 2017), Paru Co (Bird et al., 2014), and Dongge Cave (Wang et al., 2005), are indicated. (B) The coring site of Lake Kusai. Bathymetry was measured in 2002.

Sample number	Lab ID	Material	Depth (cm)	¹⁴ C age (yr BP)	Reservoir-corrected ¹⁴ C age by 3200 yr (yr BP)
KSD-1-42	NZA 50891	TOC	21	3454 ± 21	254 ± 21
KSD-1-63	NZA 50892	TOC	31.5	3611 ± 21	411 ± 21
KSD-1-96	NZA 50893	TOC	48	3614 ± 22	414 ± 22
KSD-1-133	NZA 50894	TOC	66.5	4115 ± 22	915 ± 22
KSD-1-163	NZA 50895	TOC	81.5	4196 ± 23	996 ± 23
KSD-1-190	NZA 50896	TOC	95	4656 ± 28	1456 ± 28
KSD-1-225	NZA 50897	TOC	116.5	4750 ± 23	1550 ± 23
KSD-1-251	NZA 50898	TOC	129.5	5135 ± 23	1935 ± 23
KSD-1-280	NZA 50899	TOC	144	5161 ± 23	1961 ± 23
KSD-1-318	NZA 50900	TOC	163	5248 ± 23	2048 ± 23

Table 1. AMS ¹⁴C ages and reservoir-corrected ¹⁴C ages obtained for core KSD-1.

Following the procedures above, sediment slabs were cut lengthwise and ground to a thickness of approximately 50 μ m. Optical microscope analyses were performed to count the laminations. We counted at least three times to ensure data quality and consistency. We estimated an approximate 1% error in ages based on the replicated counts.

Nondestructive XRF measurements on split long core surfaces at 0.5-cm resolution were performed with an Itrax XRF core scanner (Croudace et al., 2006) at the Institute of Geology and Geophysics, Chinese Academy of Sciences. All elements were measured at an X-ray voltage of 30 kV and a current of 55 mA. The data are semiquantitative and provide relative fluctuations in the chemical element compositions as counts per second (cps; Croudace et al., 2006; Rothwell and Rack, 2006; Löwemark et al., 2011).

RESULTS AND DISCUSSION

Varve structure

The seasonal rhythms of climate induce recurrent yearly changes in sedimentary input and varve formation (Haltia-Hovi et al., 2007; Zolitschka et al., 2015). The basic couplet structure in Lake Kusai was found to be composed of darkand light-colored lamina. The thin dark layers composed of coarse sand and silt, mainly quartz, feldspar, and mica, were formed by aeolian processes during winter. The light layers were relatively thick and consisted of authigenic carbonates and fine-grained clay minerals. These laminae were explained as representing summer season accumulations during which carbonates were deposited (Liu, X., et al., 2014b; Chen et al., 2016).

Chronology

An apparent rise in ¹³⁷Cs activity occurred at approximately 3.5 cm in the short core and can probably be explained by the onset of major fallout in AD 1954 in the Northern Hemisphere. The peak of ¹³⁷Cs at approximately 3 cm corresponds to AD 1963, which is the peak of world nuclear weapon testing (Fig. 2A; Pennington, 1973). The ¹³⁷Cs-based

chronology is generally in good agreement with the varvebased chronology but does not agree well with either ²¹⁰Pb model (CIC and CRS), an aspect which needs to be studied in the future.

The age (¹⁴C)/depth correlation for the Lake Kusai records is shown in Figure 2B. A comparison of the ¹³⁷Cs-based ages at a depth of 4 cm (AD 1950 or 0 yr BP) with the linearly fitted ¹⁴C age at the same level, which was determined by interpolation among the ten AMS ¹⁴C ages, suggests that the radiocarbon age is in excess of the 137 Cs age by 3200 yr. This phenomenon can be ascribed to the reservoir effect, which is widespread in the radiocarbon dating of lacustrine sediments in western China, particularly in saline lakes on the QTP (Herzschuh, 2006; Morrill et al., 2006; Liu, X., et al., 2008a; Hou et al., 2012). Ten AMS ¹⁴C ages from the long core have a high linear correlation ($R^2 = 0.9654$), which demonstrates that the hydrochemical conditions were stable, and the magnitude of the reservoir effect had no obvious changes. Thus, we assumed an invariable ¹⁴C reservoir effect over the time scale of the long core and subtracted a value of 3200 from all ¹⁴C ages prior to calibration. The agedepth model was constructed by the Bayesian method (Blaauw and Christen, 2011) and calibrated using the IntCal09 dataset (Reimer et al., 2013). The varve age agrees well with the calibrated ¹⁴C age, which proves that the laminations of the long core are varves and that the varve-based chronology is suitable for analyzing the decadal- to centennial-scale paleoclimatic changes of Lake Kusai. Our chronology in this study is based on the varve age. The chronology spans from approximately AD 31 to AD 2010, when the core was collected.

Temperature variations in Lake Kusai and their relationship with solar activity

Previous studies have indicated that the carbonates deposited in the sediments of Lake Kusai are predominately authigenic in origin, as a result of chemical and biochemical processes, and are mainly composed of aragonite (Yao et al., 2011). The deposition of authigenic carbonate is complex and influenced by many factors, such as temperature, salinity, and the



Figure 2. (color online) (A) Dating results of the varve and 137 Cs ages from the short core. (B) Varve and AMS 14 C ages (after 14 C ages were reservoir corrected by 3200 yr) from the KSD-1 long core.

primary productivity of lakes (Kelts and Hsü, 1978), but temperature is the most important influencing factor (Liu, X., et al., 2008b; Li and Liu, 2018). Increases in temperature can result in increased carbonate precipitation based on the following mechanisms: (1) decreases in the solubility of $CaCO_3$ (Conroy et al., 2014); (2) the evaporation of the lake region is enhanced, leading to the enrichment and supersaturation of Ca²⁺ and CO₃²⁻ in lake waters and thus facilitating the preferential precipitation of carbonates (Xiao et al., 2006; Mishra et al., 2015); (3) algae photosynthesis assimilates a high amount of CO_2 (aq) (Opitz et al., 2012), which can lead to the supersaturation of CO_3^{2-} and promote the deposition of carbonates; and (4) increases in the abundance of cyanobacteria, which serve as nuclei for carbonate precipitation (Hodell et al., 1998). Ca is mainly associated with the mineral CaCO₃ (Cohen, 2003), which is the main component of authigenic carbonates in lake sediments. Therefore, we consider temperature to be closely related to fluctuations in Ca.

To verify this link, we compared the Ca record of Lake Kusai with the δ^{18} O record from the Dunde ice core (Thompson et al., 2006) and the temperature reconstruction of the QTP based on multiple proxy records (Fig. 3; Yang et al., 2003). The fluctuations in the three curves are similar at the centennial time scale, and the small discrepancies among the three records possibly reflect errors in chronology. The Pearson correlations for the Ca record with the ice core record (R = 0.63, P < 0.01, n = 256) and the temperature reconstruction of the QTP (R = 0.22, P < 0.01, n = 363) are significant at the centennial time scale. During the last 2000 yr, three distinct high-temperature episodes that occurred in AD 31–175, 475–1400, and 1900–present correlate well with

the Roman Warm Period, Medieval Warm Period, and Current Warm Period, respectively. Between these three warm episodes are two low-temperature periods in AD 175–475 and 1400–1900, which might correspond to the Dark Age Cold Period and Little Ice Age, respectively. These warm and cold periods were also reported by other temperature records in China (Yang et al., 2002; Ge et al., 2013), which further demonstrates that the temperature fluctuations in the northern QTP were consistent with those of other areas over the last 2000 yr.

The role of solar activity on the Earth's climate system and its influence at a variety of temporal and spatial scales have been widely discussed (Beer et al., 2000; Bard and Frank, 2006; Saarni et al., 2016). Previous studies have demonstrated that the cosmogenic radionuclide ¹⁰Be can be used to reconstruct solar activity beyond the limits of instrumental records (Berggren et al., 2009; Adolphi et al., 2014). In general, variations in temperature are positively correlated with solar activity at decadal to millennial timescales (Wiles et al., 2004; Usoskin et al., 2005; Liu, Y., et al., 2009). Thus, we compared the detrended Ca record (Δ Ca) from Lake Kusai to the detrended total solar irradiance record (Δ TSI) reconstructed by ¹⁰Be (Fig. 4; Steinhilber et al., 2009). The fluctuations in Ca are generally in good agreement with the variations in TSI at decadal (R = 0.2, P < 0.01,n = 392) and centennial (R = 0.34, P < 0.01, n = 374) timescales over the last 2000 yr. The results reveal a positive correlation between solar activity and temperature variations at decadal to centennial timescales.

Furthermore, this result is also supported by cross-wavelet transform (XWT), a method for examining common periodicities and relative phases in time-frequency space between



Figure 3. Comparison of the Ca record from Lake Kusai with different paleoclimatic signals over the last 2000 yr. (A) The grey and black lines represent the decadal and centennial variations in the δ^{18} O record from the Dunde ice core, respectively (Thompson et al., 2006). (B) Ca record from Lake Kusai. The superimposed smoothed curve emphasizes the centennial long-term variations. (C) Temperature changes in the QTP reconstructed by various paleoclimatic proxy records (Yang et al., 2003).

two records (Fig. 5; Grinsted et al., 2004). We noticed that the Ca and TSI records exhibit a broad band of common frequencies extending from ~210 yr to 40 yr that are significant at a 95% confidence level relative to the red noise spectrum. More importantly, both records show a clear positive correlation with the highest power at periodicities of approximately 210 yr, which is similar to the de Vries/Suess solar activity cycle (~210 yr; Suess, 1980; Wagner et al., 2001; Huang and Tian, 2018). Our results show that the temperature fluctuations reflected by the Ca record reveal a high multi-decadal

to centennial-scale spectral coherency with solar activity. In conclusion, solar output was the likely primary forcing factor of decadal- to centennial-scale temperature changes in the northern QTP over the last 2000 yr.

Fluctuations in effective moisture over the last 2000 yr

Rb, being chemically similar to K, is often enriched in finegrained clay mineral assemblages (Koinig et al., 2003;



Figure 4. Comparison between the detrended TSI record (Δ TSI) reconstructed from the cosmogenic radionuclide ¹⁰Be data (red line; Steinhilber et al., 2009) and the detrended Ca record (Δ Ca) from Lake Kusai (green line) over the last 2000 yr. Correlations between two records are shown above the figure. The double asterisk denotes a significant correlation at the 99% confidence level. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)



Figure 5. Cross-wavelet transform for the Ca and TSI (Steinhilber et al., 2009) records over the last 2000 yr. Common spectral strengths are shown by colours ranging from deep blue (weak) to deep red (strong). The region without lighter shading delineates a range in which the identified frequencies can be interpreted. The irregular black curves delineate a 95% confidence level against the red noise signal. The small arrows illustrate the phase differences between the two data sets: the arrow pointing right is in-phase, the arrow pointing left is out-of-phase, and other angles are shown in proportion to phase differences. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Liu, X., et al., 2014a). Sr is generally a substitute for Ca in authigenic carbonates and coprecipitates within lake sediments (Cohen, 2003; Croudace and Rothwell, 2015). Sr can be dispersed in minerals containing Ca (e.g., aragonite and anorthose) with an isomorphism relationship due to their similar ionic radii. Our Sr record shows a strong positive correlation with the Ca record (R = 0.77, P < 0.01, n = 335), which demonstrates that Sr was mostly coprecipitated with Ca rather than derived from allochthonous detrital input. In general, increases in regional effective moisture lead to increases in river runoff, which can enhance the degree of catchment erosion and carry high levels of Rb into lakes with fine-grained detrital input. Thus, we used the Rb/Sr ratio to indicate the variations in effective moisture (Heymann et al., 2013) in Lake Kusai over the last 2000 yr. To avoid the closed-sum effect and show the relative fluctuations of the curve (Weltje

and Tjallingii, 2008), we used the log-ratios of Rb/Sr as the index to delineate effective moisture fluctuations in this study.

To verify the reliability of our effective moisture index, we compared the log (Rb/Sr) data from Lake Kusai to the glacial accumulation record from the Guliya ice core (Yao et al., 1996) over the last 1700 yr (Fig. 6). Over this time interval, the Pearson correlation coefficients of the two records reach high values at decadal (R = 0.44, P < 0.01, n = 337) and centennial (R = 0.66, P < 0.01, n = 319) timescales.

The stalagmite record of Dongge Cave has been widely interpreted as a proxy for the intensity and precipitation of the Indian summer monsoon (ISM) throughout the Holocene (Dykoski et al., 2005; Wang et al., 2005). To further investigate the forcing mechanism of effective moisture variations in the northern QTP, we compared the log (Rb/Sr) data of Lake Kusai with the radiometrically dated δ^{18} O record from



Figure 6. Comparison of the log (Rb/Sr) data from Lake Kusai (blue line) and the glacial accumulation record from the Guliya ice core (red line; Yao et al., 1996) over the last 1700 yr. The thin and thick lines indicate the fluctuations at decadal and centennial timescales, respectively. Correlations between the two records are shown above the figure. The double asterisk denotes a significant correlation at the 99% confidence level. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)



Figure 7. Comparison between the log-ratios of Rb/Sr from Lake Kusai (tawny line) and the δ^{18} O record from Dongge Cave (dark blue line; Wang et al., 2005) over the last 2000 yr. Correlations between the two records are shown above the figure. The double asterisk denotes a significant correlation at the 99% confidence level. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

stalagmites in Dongge Cave at the decadal to centennial timescales (Fig. 7). The results show that the effective moisture variability of Lake Kusai reflected by the log-ratios of Rb/ Sr is well-related to ISM strengthening, as indicated by the δ^{18} O record from Dongge Cave, at decadal (R = -0.26, P < 0.01, n = 392) and centennial (R = -0.45, P < 0.01, n = 374) timescales. These results imply a positive correlation between the effective moisture variations and the intensity of the ISM over the last 2000 yr, which might demonstrate that the ISM can affect not only low-latitude regions but also mid-latitude regions (the northern QTP; Liu, X., et al., 2009; Ramisch et al., 2016). Thus, our results indicate that the decadal- to centennial-scale effective moisture variations in Lake Kusai over the last 2000 yr can be mainly ascribed to the response to the ISM.

Furthermore, our result has also been evidenced by hydroclimatic records from other monsoon-influenced lakes in the southern QTP. We compared the log (Rb/Sr) data of Lake



Figure 8. Time series of log (Rb/Sr) from Lake Kusai compared with other hydroclimatic records from the southern QTP. (A) Log (Rb/Sr) record from Lake Kusai. The superimposed smoothed curve emphasizes the centennial long-term variations. (B) Paru Co PC1 grain size (Bird et al., 2014). (C) Ngamring Tso reconstructed summer (JJAS) precipitation (Conroy et al., 2017).



Figure 9. Cross-wavelet analyses of the TSI record (Steinhilber et al., 2009) with (A) the log (Rb/Sr) record and (B) the δ^{18} O record from stalagmites over the last 2000 yr (Wang et al., 2005). Common spectral strengths are shown by colours ranging from deep blue (weak) to deep red (strong). The region without lighter shading delineates a range in which the identified frequencies can be interpreted. The irregular black curves delineate a 95% confidence level against the red noise signal. The small arrows illustrate the phase differences between the two data sets: the arrow pointing right is in-phase, the arrow pointing left is out-of-phase, and other angles are shown in proportion to phase differences. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Kusai with the Paru Co lake-level curve (Bird et al., 2014) and the Ngamring Tso reconstructed summer (JJAS) precipitation (Fig. 8; Conroy et al., 2017). The fluctuations in the three curves are similar, which indicate that the moisture variability of both northern and southern parts of QTP have been driven by the ISM during the last 2000 yr. Paleoclimatic

records in the northern QTP, however, remain scarce. Hence, this conclusion needs to be further proven by more highresolution records in the future.

As an important component of the global monsoon system (Wang et al., 2017), the ISM has a profound influence on the moisture variations of East Asia. Asian monsoonal system intensity at millennial or orbital timescales is predominantly affected by changes in solar activity and insolation (Wang et al., 2005; Liu, J., et al., 2009; Cheng et al., 2016). The reliability of this assertion, however, is still under debate for short timescales. For example, as an indicator of ISM intensity, the δ^{18} O record from Dongge Cave does not agree well with the residual atmospheric ¹⁴C record (Stuiver, 1998) at the decadal time scale in the Holocene (Wang et al., 2005) and in the last 2000 yr. Thus, we performed cross-wavelet analyses of the TSI record (Steinhilber et al., 2009) with the log (Rb/Sr) record from Lake Kusai and with the δ^{18} O record from Dongge Cave (Wang et al., 2005) over the last 2000 yr (Fig. 9). Neither record revealed the evident common periodicities with solar activity at decadal to centennial timescales, even though they shared some frequencies. This result might demonstrate that the decadal- to centennial-scale effective moisture variations in the northern QTP were not directly controlled by solar activity over the last 2000 yr. This phenomenon has also been reported by other paleoclimatic studies (Yao et al., 1996; Ekdahl et al., 2008; Zhao et al., 2011; Yang et al., 2014; Conroy et al., 2017), and its underlying mechanisms remain ambiguous. At short timescales, many internal factors of climate systems, such as atmospheric circulations, large-scale ocean-atmosphere processes, and human activities, can also influence regional hydroclimatic conditions (PAGES, 2009; Bird et al., 2016; Conroy et al., 2017; PAGES Hydro2k Consortium, 2017), and this topic still needs to be studied in the future.

CONCLUSIONS

Based on the high-resolution XRF scanning records from 1.52-m-long succession of varved sediments from Lake Kusai, we draw the following major conclusions:

- (1) At decadal to centennial timescales, the Ca record is a proxy for temperature changes, which correlates well with the variations and periodicities of solar activity. These results indicate that solar output was the likely primary forcing mechanism of decadal- to centennialscale temperature fluctuations over the last 2000 yr.
- (2) The decadal- to centennial-scale variability of the effective moisture of Lake Kusai, which is reflected by the log-ratios of Rb/Sr, was mainly influenced by the ISM over the last 2000 yr. However, we have not found obvious periodicities of solar activity in our effective moisture record. This finding indicates that the evolution of decadal- to centennial-scale effective moisture in the northern QTP might not have been directly controlled by solar activity over the last 2000 yr.

ACKNOWLEDGMENTS

We would like to express our gratitude to senior editor Lewis Owen, associate editor Peter Langdon, Professor Jonathan Holmes, and an anonymous reviewer for their constructive comments which improved the quality of this manuscript. We thank Kai Li (Zhejiang Normal University), Junfeng Li (Capital Normal University), and Zhenyu Ni (Nanjing Institute of Geography and Limnology, Chinese Academy of Sciences) for their help with the fieldwork. We also acknowledge American Journal Experts for the Englishlanguage editing. This study was supported by the National Natural Science Foundation of China (NSFC) and the German Research Foundation (DFG; Grant No. 41861134030), and the NSFC (Grant No. 41572338).

REFERENCES

- Adolphi, F., Muscheler, R., Svensson, A., Aldahan, A., Possnert, G., Beer, J., Sjolte, J., Björck, S., Matthes, K., Thiéblemont, R., 2014. Persistent link between solar activity and Greenland climate during the Last Glacial Maximum. *Nature Geoscience* 7, 662–666.
- An, Z.S., Colman, S.M., Zhou, W.J., Li, X.Q., Brown, E.T., Jull, A.J.T., Cai, Y.J., *et al.*, 2012. Interplay between the Westerlies and Asian monsoon recorded in Lake Qinghai sediments since 32 ka. *Scientific Reports* 2, 619.
- An, Z.S., Kutzbach, J.E., Prell, W.L., Porter, S.C., 2001. Evolution of Asian monsoons and phased uplift of the Himalaya-Tibetan plateau since Late Miocene times. *Nature* 411, 62–66.
- Appleby, P.G., 2001. Chronostratigraphic Techniques in Recent Sediments. In: Last, W.M. and Smol, J.P. (Eds.), *Tracking Environmental Change Using Lake Sediments. Volume 1: Basin Analysis, Coring, and Chronological Techniques.* Kluwer Academic Publishers, Dordrecht, pp. 171–203.
- Appleby, P.G., Oldfieldz, F., 1983. The assessment of ²¹⁰Pb data from sites with varying sediment accumulation rates. *Hydrobiolo*gia 103, 29–35.
- Bard, E., Frank, M., 2006. Climate change and solar variability: What's new under the sun? *Earth and Planetary Science Letters* 248, 1–14.
- Beer, J., Mende, W., Stellmacher, R., 2000. The role of the sun in climate forcing. *Quaternary Science Reviews* 19, 403–415.
- Berggren, A.M., Beer, J., Possnert, G., Aldahan, A., Kubik, P., Christl, M., Johnsen, S.J., Abreu, J., Vinther, B.M., 2009. A 600-year annual ¹⁰Be record from the NGRIP ice core, Greenland. *Geophysical Research Letters* 36, 431–433.
- Bird, B.W., Lei, Y.B., Perello, M., Polissar, P.J., Yao, T.D., Finney, B., Bain, D., Pompeani, D., Thompson, L.G., 2016. Late-Holocene Indian summer monsoon variability revealed from a 3300-year-long lake sediment record from Nir'pa Co, southeastern Tibet. *Holocene* 27, 1–12.
- Bird, B.W., Polisar, P.J., Lei, Y.B., Thompson, L.G., Yao, T.D., Finney, B.P., Bain, D.J., Pompeani, D.P., Steinman, B.A., 2014. A Tibetan lake sediment record of Holocene Indian summer monsoon variability. *Earth and Planetary Science Letters* 399, 92–102.
- Blaauw, M., Christen, J.A., 2011. Flexible paleoclimate age-depth models using an autoregressive gamma process. *Bayesian Analy*sis 6, 457–474.
- Chen, Y., Liu, X.Q., He, L., Ye, L., Chen, H.F., Li, K., 2016. Microarea analysis and mechanism of varves from Lake Kusai in the Hoh Xil area, northern Tibetan Plateau. [In Chinese with English abstract.] *Acta Geologica Sinica* 90, 1006–1015.
- Cheng, H., Edwards, R.L., Sinha, A., Spötl, C., Yi, L., Chen, S.T., Kelly, M., *et al.*, 2016. The Asian monsoon over the past 640,000 years and ice age terminations. *Nature* 534, 640.
- Cohen, A.S., 2003. Paleolimnology: The History and Evolution of Lake Systems. Oxford University Press, Oxford.

- Conroy, J.L., Hudson, A.M., Overpeck, J.T., Liu, K.B., Wang, L., Cole, J.E., 2017. The primacy of multidecadal to centennial variability over late-Holocene forced change of the Asian Monsoon on the southern Tibetan Plateau. *Earth and Planetary Science Letters* 458, 337–348.
- Conroy, J.L., Thompson, D.M., Collins, A., Overpeck, J.T., Bush, M.B., Cole, J.E., 2014. Climate influences on water and sediment properties of Genovesa Crater Lake, Galápagos. *Journal* of Paleolimnology 52, 331–347.
- Croudace, I.W., Rindby, A., Rothwell, R.G., 2006. ITRAX: description and evaluation of a new multi-function X-ray core scanner. In: Rothwell, R.G. (Ed.), *New Techniques in Sediment Core Analysis*. Geological Society, London, pp. 51–63.
- Croudace, I.W., Rothwell, R.G., 2015. Micro-XRF Studies of Sediment Cores: Applications of a Non-Destructive Tool for the Environmental Sciences. Springer, Dordrecht.
- Dykoski, C.A., Edwards, R.L., Cheng, H., Yuan, D.X., Cai, Y.J., Zhang, M.L., Lin, Y.S., Qing, J.M., An, Z.S., Revenaugh, J., 2005. A high-resolution, absolute-dated Holocene and deglacial Asian monsoon record from Dongge Cave, China. *Earth and Planetary Science Letters* 233, 71–86.
- Ekdahl, E.J., Fritz, S.C., Baker, P.A., Rigsby, C.A., Coley, K., 2008. Holocene multi-decadal to millennial-scale hydrologic variability on the South American Altiplano. *Holocene* 18, 867.
- Fang, Z.Y., Zhu, F.K., Jiang, J.X., Qian, Z.A., 1997. Research of Dust-storm in China. [In Chinese.] Meteorological Press, Beijing.
- Gasse, F., Arnold, M., Fontes, J.C., Fort, M., Gibert, E., Huc, A., Li, B., Li, Y., Liu, Q., Mélières, F., 1991. A 13,000-year climate record from western Tibet. *Nature* 353, 742–745.
- Ge, Q.S., Zheng, J.Y., Hao, Z.X., Liu, H.L., 2013. General characteristics of climate changes during the past 2000 years in China. *Science in China Series D Earth Sciences* 56, 321–329.
- Grießinger, J., Bräuning, A., Helle, G., Hochreuther, P., Schleser, G., 2017. Late Holocene relative humidity history on the southeastern Tibetan plateau inferred from a tree-ring δ ¹⁸O record: Recent decrease and conditions during the last 1500 years. *Quaternary International* 430, 52–59.
- Grinsted, A., Moore, J.C., Jevrejeva, S., 2004. Application of the cross wavelet transform and wavelet coherence to geophysical time series. *Nonlinear Processes in Geophysics* 11, 561–566.
- Haltia-Hovi, E., Saarinen, T., Kukkonen, M., 2007. A 2000-year record of solar forcing on varved lake sediment in eastern Finland. *Quaternary Science Reviews* 26, 678–689.
- Herzschuh, U., 2006. Palaeo-moisture evolution in monsoonal Central Asia during the last 50,000 years. *Quaternary Science Reviews* 25, 163–178.
- Herzschuh, U., Borkowski, J., Schewe, J., Mischke, S., Tian, F., 2014. Moisture-advection feedback supports strong early-to-mid Holocene monsoon climate on the eastern Tibetan Plateau as inferred from a pollen-based reconstruction. *Palaeogeography*, *Palaeoclimatology*, *Palaeoecology* 402, 44–54.
- Heymann, C., Nelle, O., Dörfler, W., Zagana, H., Nowaczyk, N., Xue, J.B., Unkel, I., 2013. Late Glacial to mid-Holocene palaeoclimate development of Southern Greece inferred from the sediment sequence of Lake Stymphalia (NE-Peloponnese). *Quaternary International* 302, 42–60.
- Hodell, D.A., Schelske, C.L., Fahnenstiel, G.L., Robbins, L.L., 1998. Biologically induced calcite and its isotopic composition in Lake Ontario. *Limnology and Oceanography* 43, 187–199.
- Hou, J.Z., D'Andrea, W.J., Liu, Z.H., 2012. The influence of ¹⁴C reservoir age on interpretation of paleolimnological records from the Tibetan Plateau. *Quaternary Science Reviews* 48, 67–79.

- Huang, C.J., Tian, X.L., 2018. Discussion on the driving mechanism of solar activity to interannual-suborbital-scale climate change. [In Chinese with English abstract.] *Quaternary Science* 38, 1255–1267.
- Kelts, K., Hsü, K.J., 1978. Freshwater Carbonate Sedimentation. In: Lerman, A. (Ed.), *Lakes: Chemistry, Geology, Physics*. Springer, New York, pp. 295–323.
- Koinig, K.A., Shotyk, W., Lotter, A.F., Ohlendorf, C., Sturm, M., 2003. 9000 years of geochemical evolution of lithogenic major and trace elements in the sediment of an alpine lake-the role of climate, vegetation, and land-use history. *Journal of Paleolimnology* 30, 307–320.
- Li, B.Y., 1996. Environment of Kekexili area, Qinghai Province. [In Chinese.] Science Press, Beijing.
- Li, J.F., Liu, X.Q., 2018. Orbital- and suborbital-scale changes in the East Asian summer monsoon since the last deglaciation. *Holocene* 28, 1216–1224.
- Li, K., Liu, X.Q., Wang, Y.B., Herzschuh, U., Ni, J., Liao, M.N., Xiao, X.Y., 2017. Late Holocene vegetation and climate change on the southeastern Tibetan Plateau: Implications for the Indian Summer Monsoon and links to the Indian Ocean Dipole. *Quaternary Science Reviews* 177, 235–245.
- Liu, J., Wang, B., Ding, Q.H., Kuang, X.Y., Willie, S., Eduardo, Z., 2009. Centennial variations of the global monsoon precipitation in the last millennium: Results from ECHO-G model. *Journal of Climate* 22, 2356–2371.
- Liu, X.Q., Dong, H.L., Rech, J.A., Matsumoto, R., Bo, Y., Wang, Y.B., 2008a. Evolution of Chaka Salt Lake in NW China in response to climatic change during the Latest Pleistocene-Holocene. *Quaternary Science Reviews* 27, 867–879.
- Liu, X.Q., Dong, H.L., Yang, X.D., Herzschuh, U., Zhang, E.L., Stuut, J.B.W., Wang, Y.B., 2009. Late Holocene forcing of the Asian winter and summer monsoon as evidenced by proxy records from the northern Qinghai-Tibetan Plateau. *Earth and Planetary Science Letters* 280, 276–284.
- Liu, X.Q., Herzschuh, U., Shen, J., Jiang, Q.F., Xiao, X.Y., 2008b. Holocene environmental and climatic changes inferred from Wulungu Lake in northern Xinjiang, China. *Quaternary Research* 70, 412–425.
- Liu, X.Q., Herzschuh, U., Wang, Y.B., Kuhn, G., Yu, Z.T., 2014a. Glacier fluctuations of Muztagh Ata and temperature changes during the late Holocene in westernmost Tibetan Plateau, based on glaciolacustrine sediment records. *Geophysical Research Letters* 41, 6265–6273.
- Liu, X.Q., Yu, Z.T., Dong, H.L., Chen, H.F., 2014b. A less or more dusty future in the Northern Qinghai-Tibetan Plateau? *Scientific Reports* 4, 6672.
- Liu, Y., An, Z.S., Linderholm, H.W., Chen, D.L., Song, H.M., Cai, Q.F., Sun, J.Y., Hua, T., 2009. Annual temperatures during the last 2485 years in the mid-eastern Tibetan Plateau inferred from tree rings. *Science in China Series D Earth Sciences* 52, 348–359.
- Löwemark, L., Chen, H.F., Yang, T.N., Kylander, M., Yu, E.F., Hsu, Y.W., Lee, T.Q., Song, S.R., Jarvis, S., 2011. Normalizing XRF-scanner data: A cautionary note on the interpretation of high-resolution records from organic-rich lakes. *Journal of Asian Earth Sciences* 40, 1250–1256.
- Mishra, P.K., Anoop, A., Schettler, G., Prasad, S., Jehangir, A., Menzel, P., Naumann, R., Yousuf, A.R., Basavaiah, N., Deenadayalan, K., 2015. Reconstructed late Quaternary hydrological changes from Lake Tso Moriri, NW Himalaya. *Quaternary International* 371, 76–86.

- Morrill, C., Overpeck, J.T., Cole, J.E., Liu, K.B., Shen, C., Tang, L., 2006. Holocene variations in the Asian monsoon inferred from the geochemistry of lake sediments in central Tibet. *Quaternary Research* 65, 232–243.
- Oldfield, F., Appleby, P.G., 1984. Empirical testing of ²¹⁰Pb-dating models for lake sediments. In: Haworth, E.Y., Lund, J.W.G. (Eds.), *Lake Sediments and Environmental History*. Leicester University Press, Leicester, pp. 93–124.
- Opitz, S., Wünnemann, B., Aichner, B., Dietze, E., Hartmann, K., Herzschuh, U., Ijmker, J., *et al.*, 2012. Late Glacial and Holocene development of Lake Donggi Cona, north-eastern Tibetan Plateau, inferred from sedimentological analysis. *Palaeogeography, Palaeoclimatology, Palaeoecology* 337–338, 159–176.
- Past Global Changes (PAGES), 2009. Past Global Changes: Science Plan and Implementation Strategy (IGBP Report No. 57). IGBP Secretariat, Stockholm.
- Past Global Changes (PAGES) Hydro2k Consortium, 2017. Comparing proxy and model estimates of hydroclimate variability and change over the Common Era. *Climate of the Past* 13, 1851–1900.
- Pennington, W., 1973. The recent sediments of Windermere. *Freshwater Biology* 3, 363–382.
- Ramisch, A., Lockot, G., Haberzettl, T., Hartmann, K., Kuhn, G., Lehmkuhl, F., Schimpf, S., *et al.*, 2016. A persistent northern boundary of Indian Summer Monsoon precipitation over Central Asia during the Holocene. *Scientific Reports* 6, 25791.
- Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey, C.B., Buck, C.E., *et al.*, 2013. IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP. *Radiocarbon* 51, 1111–1150.
- Rothwell, R.G., Rack, F.R., 2006. New techniques in sediment core analysis: an introduction. In: Rothwell, R.G. (Ed.), *New Techniques in Sediment Core Analysis*. Special Publication, Vol. 267. Geological Society, London, pp. 1–29.
- Saarni, S., Muschitiello, F., Weege, S., Brauer, A., Saarinen, T., 2016. A late Holocene record of solar-forced atmospheric blocking variability over Northern Europe inferred from varved lake sediments of Lake Kuninkaisenlampi. *Quaternary Science Reviews* 154, 100–110.
- Shao, X.M., Huang, L., Liu, H.B., Liang, E.Y., Fang, X.Q., Wang, L.L., 2005. Reconstruction of precipitation variation from tree rings in recent 1000 years in Delingha, Qinghai. *Science in China Series D Earth Sciences* 48, 939–949.
- Shen, J., Liu, X.Q., Wang, S.M., Matsumoto, R., 2005. Palaeoclimatic changes in the Qinghai Lake area during the last 18,000 years. *Quaternary International* 136, 131–140.
- Steinhilber, F., Beer, J., Fröhlich, C., 2009. Total solar irradiance during the Holocene. *Geophysical Research Letters* 36, 308–308.
- Stuiver, M., 1998. INTCAL 98 radiocarbon age calibration 24,000 cal BP. *Radiocarbon* 40, 1041–1083.
- Suess, H.E., 1980. The radiocarbon record in tree rings of the last 8000 years. *Radiocarbon* 22, 200–209.
- Tan, L.C., Cai, Y.J., An, Z.S., Edwards, R.L., Cheng, H., Shen, C.C., Zhang, H.W., 2010. Centennial to decadal-scale monsoon precipitation variability in the semi-humid region, northern China during the last 1860 years: Records from stalagmites in Huangye Cave. *Holocene* 21, 287–296.
- Tan, L.C., Cai, Y.J., Yi, L., An, Z.S., Ai, L., 2008. Precipitation variations of Longxi, northeast margin of Tibetan Plateau since AD 960 and their relationship with solar activity. *Climate of the Past* 20, 19–28.

- Thompson, L.G., Mosleythompson, E., Brecher, H., Davis, M., León, B., Les, D., Lin, P.N., Mashiotta, T., Mountain, K., 2006. Abrupt tropical climate change: Past and present. *Proceedings of the National Academy of Sciences of the United States of America* 103, 10536–10543.
- Thompson, L.G., Yao, T.D., Mosley-Thompson, E., Davis, M.E., Henderson, K.A., Lin, P.N., 2000. A high-resolution millennial record of the south asian monsoon from Himalayan ice cores. *Science* 289, 1916–1920.
- Usoskin, I.G., Schüssler, M., Solanki, S.K., Mursula, K., 2005. Solar activity, cosmic rays, and Earth's temperature: A millennium-scale comparison. *Journal of Geophysical Research* 110, A10102.
- Wagner, G., Beer, J., Masarik, J., Muscheler, R., Kubik, P.W., Mende, W., Laj, C., Raisbeck, G.M., Yiou, F., 2001. Presence of the Solar de Vries Cycle (~205 years) during the Last Ice Age. *Geophysical Research Letters* 28, 303–306.
- Wang, P.X., Wang, B., Cheng, H., Fasullo, J., Guo, Z., Kiefer, T., Liu, Z.Y., 2017. The global monsoon across time scales: Mechanisms and outstanding issues. *Earth-Science Reviews* 174, 84–121.
- Wang, S.M., Dou, H.S., 1998. *Lakes in China*. [In Chinese.] Science Press, Beijing.
- Wang, Y.J., Cheng, H., Edwards, R.L., He, Y.Q., Kong, X.G., An, Z.S., Wu, J.Y., Kelly, M.J., Dykoski, C.A., Li, X.D., 2005. The Holocene Asian monsoon: Links to solar changes and North Atlantic climate. *Science* 308, 854–857.
- Weltje, G.J., Tjallingii, R., 2008. Calibration of XRF core scanners for quantitative geochemical logging of sediment cores: Theory and application. *Earth and Planetary Science Letters* 274, 423–438.
- Wiles, G.C., D'Arrigo, R.D., Villalba, R., Calkin, P.E., Barclay, D.J., 2004. Century-scale solar variability and Alaskan temperature change over the past millennium. *Geophysical Research Letters* 31, 305–316.
- Xiao, J.L., Wu, J.T., Si, B., Liang, W.D., Nakamura, T., Liu, B.L., Inouchi, Y., 2006. Holocene climate changes in the monsoon/ arid transition reflected by carbon concentration in Daihai Lake of Inner Mongolia. *Holocene* 16, 551–560.
- Yan, H., Soon, W., Wang, Y.H., 2015. A composite sea surface temperature record of the northern South China Sea for the past 2500 years: A unique look into seasonality and seasonal climate changes during warm and cold periods. *Earth-Science Reviews* 141, 122–135.
- Yang, B., Braeuning, A., Johnson, K.R., Shi, Y., 2002. General characteristics of temperature variation in China during the last two millennia. *Geophysical Research Letters* 29, 38-31–38-34.
- Yang, B., Braeuning, A., Shi, Y.F., 2003. Late Holocene temperature fluctuations on the Tibetan Plateau. *Quaternary Science Reviews* 22, 2335–2344.
- Yang, B., Qin, C., Wang, J.L., He, M.H., Melvin, T.M., Osborn, T.J., Briffa, K.R., 2014. A 3,500-year tree-ring record of annual precipitation on the northeastern Tibetan Plateau. *Proceedings of the National Academy of Sciences of the United States* of America 111, 2903–2908.
- Yao, B., Liu, X.Q., Wang, Y.B., Yang, B., 2011. Late Holocene climatic changes revealed by mineralogical records from lacustrine core KS-2006 from Lake Kusai in the Hoh Xil area, northern Tibetan Plateau. [In Chinese with English abstract.] *Journal of Lake Sciences* 23, 903–909.
- Yao, T.D., Masson-Delmotte, V., Gao, J., Yu, W.S., Yang, X.X., Risi, C., Sturm, C., *et al.*, 2013. A review of climatic controls

on δ^{18} O in precipitation over the Tibetan Plateau: Observations and simulations. *Reviews of Geophysics* 51, 525–548.

- Yao, T.D., Thompson, L.G., Qing, D.H., Tian, L.D., Jiao, K.Q., Yang, Z.H., Xie, C., 1996. Variations in temperature and precipitation in the past 2000 a on the Xizang (Tibet) Plateau-Guliya ice core record. *Science in China Series D Earth Sciences* 39, 425–433.
- Zhang, P.Z., Cheng, H., Edwards, R.L., Chen, F.H., Wang, Y.J., Yang, X.L., Liu, J., *et al.*, 2008. A test of climate, sun, and culture relationships from an 1810-year Chinese cave record. *Science* 322, 940–942.
- Zhao, P., Zhou, X.J., Liu, G., 2011. Decadal-centennial-scale change in Asian-Pacific summer thermal contrast and solar activity. *Chinese Science Bulletin* 56, 3012–3018.
- Zhu, L.P., Wu, Y.H., Wang, J.B., Xiao, L., Ju, J.T., Xie, M.P., Li, M.H., Mäusbacher, R., Schwalb, A., Daut, G., 2008. Environmental changes since 8.4 ka reflected in the lacustrine core sediments from Nam Co, central Tibetan Plateau, China. *Holocene* 18, 831–839.
- Zolitschka, B., Francus, P., Ojala, A.E.K., Schimmelmann, A., 2015. Varves in lake sediments—A review. *Quaternary Science Reviews* 117, 1–41.