


# A process-based evaluation of biases in extratropical stratosphere-troposphere coupling in subseasonal forecast systems



Zachary Lawrence, Chaim I. Garfinkel, Amy Butler, Etienne Dunn-Sigouin, Irina Statnaia, Alexey Karpechko, Gerbrand Koren, Marta Abalos, Blanca Ayarzaguen, David Barriopedro, Natalia Calvo, Alvaro de la Camara, Andrew Charlton-Perez, Daniela I.V. Domeisen, Javier Garcia-Serrano, Neil P. Hindley, Martin Jucker, Hera Kim, Simon H. Lee, Marisol Osman, Froila M. Palmeiro, Inna Polichtchouk, Jian Rao, Jadwiga H. Richter, Chen Schwartz, Seok-Woo Son, Masakazu Taguchi, Nicholas L. Tyrrell, Corwin J. Wright, Rachel W.-Y. Wu

## Introduction

- Two-way coupling between the stratosphere and troposphere is recognized as an important source of subseasonal-to-seasonal (S2S) predictability and can provide forecast windows of opportunity.
- Model biases can, however, lead to a poor representation of such coupling processes; at lead times of one to two weeks, drifts in a model's circulation related to model biases, resolution, and parameterizations have the potential to feed back on the circulation and affect stratosphere-troposphere coupling.

## Conclusions

### Northern Hemisphere:


- Nearly all S2S forecast systems underestimate the strength of the observed upward coupling from the troposphere to the stratosphere and downward coupling within the stratosphere.
- While downward coupling from the lower stratosphere to the near surface is well represented in the multi-model ensemble mean, there is substantial inter-model spread. This is likely related to overly fast decay of simulated lower stratospheric temperature anomalies.

### Southern Hemisphere:


- The forecast systems overestimate the upward coupling of wave-1 from the troposphere to the lower stratosphere, and the stratospheric vortex is over-sensitive to upward propagating wave flux.
- Forecast systems generally overestimate the strength of downward coupling from the lower stratosphere to the troposphere, even as they underestimate the radiative persistence in the lower stratosphere.

In both hemispheres, models with higher lids and a better representation of tropospheric quasi-stationary waves generally perform better at simulating these coupling processes.


## 2. Upward coupling: too-weak penetration of wv1




### What explains intermodel spread in the regression coefficients?




## 3. Summary of biases in coupling strength



## 4. Too-weak sensitivity of polar vortex to 100hPa heat flux



## 5. Too-strong sensitivity of 850hPa NAM to lower strat



## 6. Underestimated interannual variability in wave extremes

