

# Supplementary information for EGU25-11448 Peatland shrub roots increase resource acquisition with warming

Tiia Määttä<sup>1</sup>, Nikhil Chari<sup>2,3</sup>, Joanne Childs<sup>4,5</sup>, Colleen Iversen<sup>4,5</sup>, Verity Salmon<sup>4,5</sup>, Geoffrey Schwaner<sup>4,5</sup>, Sören Weber<sup>4,5</sup>, and Avni Malhotra<sup>1,6</sup>

<sup>1</sup> Department of Geography, Faculty of Science, University of Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland

<sup>2</sup> Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA, USA

<sup>3</sup> Arnold Arboretum of Harvard University, Boston, MA, USA

<sup>4</sup> Environmental Sciences Division, Oak Ridge National Laboratory, Oak Ridge, TN 37830, USA

<sup>5</sup> Climate Change Science Institute, Oak Ridge National Laboratory, Oak Ridge, TN 37830, USA

<sup>6</sup> Biological Sciences Division, Pacific Northwest National Laboratory, 902 Battelle Boulevard, Richland, WA, USA

**Correspondence to: Tiia Määttä (tiia.maatta@geo.uzh.ch)**

**1. Detailed linear mixed effects model (fitted with restricted maximum likelihood) results shown in the poster. Significant ( $p < 0.05$ ) terms are highlighted with an asterisk. All models shown here are the final models after backward variable selection with log likelihood ratio tests. See full models in table 2. Note that in all, except tree root C:N, models, the trait (response variable) was log-transformed to meet normality assumptions. The  $R^2$  values shown in the poster are marginal  $R^2$ . In all models, the reference level for  $CO_2$  treatment was “ambient”. SE = standard error, SRL = specific root length ( $m\ g^{-1}$ ), RTD = root tissue density ( $g\ cm^{-3}$ ), soil temp = soil temperature ( $^{\circ}C$ ), soil moist = soil moisture (g/g),  $eCO_2$ =elevated  $CO_2$  treatment. Unit of fine root length production is  $km\ m^{-2}$  growing season<sup>-1</sup>.**

Model	<i>p</i> -value	Slope	SE	Marginal $R^2$	Conditional $R^2$	Degrees of freedom	Random variation explained (%)
<b><u>Fine root length production</u></b>							
<b>Shrub model</b>				0.11	0.54		48.3
- Intercept	0	8.076	0.291			22	
- Soil temp	0.077	0.085	0.046			22	

- eCO <sub>2</sub>	0.358	0.185	0.197			22	
- Soil temp: eCO <sub>2</sub>	0.065	-0.096	0.049			22	
<b>Tree model</b>				0.1	0.67		62.8
- Intercept	0	7.043	0.552			24	
- Soil moist*	0.024	-0.542	0.224			24	
<b><u>SRL</u></b>							
<b>Shrub model</b>				0.25	0.51		35.1
- Intercept	0	5.893	0.09			23	
- Soil temp*	0.027	0.034	0.014			23	
- eCO <sub>2</sub>	0.054	-0.162	0.08			23	
<b>Tree model</b>				0.09	0.15		5.7
- Intercept	0	3.36	0.083			24	
- Soil temp	0.123	-0.031	0.019			24	
<b><u>Root C:N</u></b>							
<b>Shrub model</b>				0.34	0.48		21.5
- Intercept	0	3.744	0.068			12	
- Soil temp*	0.029	-0.033	0.013			12	
<b>Tree model</b>				0.03	0.4		21.5
- Intercept	0	38.218	3.745			12	
- eCO <sub>2</sub>	0.442	2.569	3.231			12	
<b><u>RTD</u></b>							
<b>Shrub model</b>				0.07	0.74		72.7
- Intercept	0	-1.352	0.207			19	
- Soil temp	0.183	-0.03	0.022			19	
<b>Tree model</b>				0.2	0.6		50.4
- Intercept	0	-1.277	0.133			17	
- Soil temp	0.82	-0.004	0.015			17	
- eCO <sub>2</sub>	0.057	-0.181	0.088			17	
- Soil temp:eCO <sub>2</sub>	0.045	-0.04	0.018			17	

**2. Full linear mixed effects model (fitted with restricted maximum likelihood) results. Significant ( $p < 0.05$ ) terms are highlighted with an asterisk. Note that in all, except tree C:N, models, the trait (response variable) was log-transformed to meet normality assumptions. In all models, the reference level for CO<sub>2</sub>**

treatment was “ambient”. SE = standard error, SRL = specific root length ( $\text{m g}^{-1}$ ), RTD = root tissue density ( $\text{g cm}^{-3}$ ), soil temp = soil temperature ( $^{\circ}\text{C}$ ), soil moist = soil moisture ( $\text{g/g}$ ),  $\text{eCO}_2$  = elevated  $\text{CO}_2$  treatment. Unit of fine root length production is  $\text{km m}^{-2}$  growing season $^{-1}$ .

Model	p-value	Slope	SE	Marginal $R^2$	Conditional $R^2$	Degrees of freedom	Random variation explained (%)
<b><u>Fine root length production</u></b>							
<b>Shrub model</b>				0.13	0.41		32.1
- Intercept	0	8.094	0.251			21	
- Soil temp	0.171	0.067	0.047			21	
- Soil moist	0.295	-0.163	0.152			21	
- $\text{eCO}_2$	0.509	0.141	0.21			21	
- Soil temp: $\text{eCO}_2$	0.056	-0.106	0.052			21	
<b>Tree model</b>				0.1	0.76		73.2
- Intercept	0	7.242	0.697			21	
- Soil temp	0.455	0.065	0.086			21	
- Soil moist	0.069	-0.492	0.257			21	
- $\text{eCO}_2$	0.232	-0.421	0.342			21	
- Soil temp: $\text{eCO}_2$	0.597	0.043	0.08			21	
<b><u>SRL</u></b>							
<b>Shrub model</b>				0.25	0.52		36.7
- Intercept	0	5.89	0.096			21	
- Soil temp	0.101	0.031	0.018			21	
- Soil moist	0.913	0.007	0.059			21	
- $\text{eCO}_2$	0.077	-0.16	0.086			21	
- Soil temp: $\text{eCO}_2$	0.666	0.009	0.02			21	
<b>Tree model</b>				0.16	0.26		11.6
- Intercept	0	3.334	0.12			21	
- Soil temp	0.211	-0.035	0.027			21	
- Soil moist	0.111	0.149	0.089			21	
- $\text{eCO}_2$	0.772	0.043	0.147			21	
- Soil temp: $\text{eCO}_2$	0.649	0.016	0.035			21	

---

**Root C:N**


<b>Shrub model</b>				0.4	0.51		19.5
- Intercept	0	3.703	0.086			9	
- Soil temp	0.065	-0.035	0.016			9	
- Soil moist	0.453	-0.044	0.055			9	
- eCO <sub>2</sub>	0.559	0.062	0.102			9	
- Soil temp: eCO <sub>2</sub>	0.956	-0.001	0.023			9	
<b>Tree model</b>				0.04	0.47		44.8
- Intercept	0	37.057	4.771			9	
- Soil temp	0.653	-0.365	0.784			9	
- Soil moist	0.711	0.991	2.592			9	
- eCO <sub>2</sub>	0.322	4.651	4.44			9	
- Soil temp: eCO <sub>2</sub>	0.452	0.639	0.813			9	

---

**RTD**

<b>Shrub model</b>				0.12	0.74		70.2
- Intercept	0	-1.384	0.217			16	
- Soil temp	0.225	-0.035	0.028			16	
- Soil moist	0.633	-0.041	0.084			16	
- eCO <sub>2</sub>	0.572	0.068	0.117			16	
- Soil temp: eCO <sub>2</sub>	0.738	-0.009	0.027			16	
<b>Tree model</b>				0.17	0.64		57
- Intercept	0	-1.31	0.136			16	
- Soil temp	0.857	0.003	0.018			16	
- Soil moist	0.198	0.068	0.051			16	
- eCO <sub>2</sub>	0.202	-0.127	0.095			16	
- Soil temp: eCO <sub>2</sub>	0.062	-0.04	0.02			16	

**3. Shrub and tree root traits with year of sampling shown in different colors.**  
**Note: year was a random effect in the linear mixed effects models. Note also the different y axis scales between shrub and tree plots.**



4. Linear regression (with year as a fixed effect and without random effects or variance structures) results. All the results are based on “full” models without backward variable selection. Significant ( $p < 0.05$ ) terms are highlighted with an asterisk. Note that in all, except root C:N and shrub RTD, models, the trait (response variable) was log-transformed to meet normality assumptions, and that interactions with year and the treatments were not included due to the small sample size. In all models, the reference level for CO<sub>2</sub> treatment was “ambient” and for year “2014” (the beginning of the SPRUCE experiment). SE = standard error, SRL = specific root length (m g<sup>-1</sup>), RTD = root tissue density (g cm<sup>-3</sup>), soil temp = soil temperature (°C), soil moist = soil moisture (g/g), eCO<sub>2</sub>=elevated CO<sub>2</sub> treatment. Unit of fine root length production is km m<sup>-2</sup> growing season<sup>-1</sup>.

Model	<i>p</i> -value	Slope	SE	Adjusted R <sup>2</sup>	Model <i>p</i> -value
<b><u>Fine root length production</u></b>					
<b>Shrub model</b>				0.15	0.174
- Intercept	<0.001	7.343	0.636		
- Soil temp	0.214	0.129	0.1		
- Soil moist	0.83	0.067	0.309		
- eCO <sub>2</sub>	0.439	0.297	0.377		
- Year					
- 2015	0.816	-0.139	0.59		
- 2016	0.291	1.133	1.046		
- 2017	0.166	1.028	0.716		
- 2022	0.055	2.021	0.995		
- Soil temp: eCO <sub>2</sub>	0.437	-0.069	0.087		
<b>Tree model</b>				0.66	<0.001
- Intercept	<0.001	6.063	0.545		
- Soil temp	0.189	0.126	0.093		
- Soil moist	0.179	-0.375	0.27		
- eCO <sub>2</sub>	0.214	-0.435	0.34		
- Year					
- 2015	0.988	-0.008	0.529		
- 2016	0.17	1.311	0.922		
- 2017	0.348	0.596	0.62		
- 2022*	0.0002	4.029	0.884		
- Soil temp: eCO <sub>2</sub>	0.566	0.047	0.08		
<b><u>SRL</u></b>					
<b>Shrub model</b>				0.36	0.021
- Intercept	<0.001	5.475	0.143		
- Soil temp*	0.027	0.054	0.023		
- Soil moist	0.262	0.08	0.069		

---

- eCO <sub>2</sub>	0.097	-0.147	0.085		
- Year					
- 2015*	0.018	0.34	0.133		
- 2016*	0.026	0.563	0.235		
- 2017*	0.014	0.434	0.161		
- 2022*	0.005	0.707	0.224		
- Soil temp: eCO <sub>2</sub>	0.605	0.01	0.02		
<b>Tree model</b>				0.1	0.251
- Intercept	<0.001	2.962	0.241		
- Soil temp	0.701	-0.016	0.041		
- Soil moist	0.076	0.223	0.119		
- eCO <sub>2</sub>	0.708	0.057	0.15		
- Year					
- 2015	0.1	0.406	0.234		
- 2016	0.39	0.358	0.408		
- 2017	0.127	0.436	0.275		
- 2022	0.122	0.631	0.391		
- Soil temp: eCO <sub>2</sub>	0.684	0.015	0.035		

---

### Root C:N

<b>Shrub model</b>				0.38	0.101
- Intercept	<0.001	42.208	5.219		
- Soil temp	0.101	-1.897	1.039		
- Soil moist	0.646	-1.442	3.032		
- eCO <sub>2</sub>	0.444	3.94	4.917		
- Year					
- 2015	-	-	-		
- 2016	0.837	1.92	9.079		
- 2017	-	-	-		
- 2022	0.363	-8.711	9.098		
- Soil temp: eCO <sub>2</sub>	0.924	-0.108	-0.098		
<b>Tree model</b>				0.17	0.281
- Intercept	<0.001	41.277	6.029		
- Soil temp	0.47	-0.771	1.022		
- Soil moist	0.793	0.765	2.829		
- eCO <sub>2</sub>	0.291	5.086	4.533		
- Year					
- 2015	-	-	-		
- 2016	0.864	-1.737	9.833		
- 2017	-	-	-		
- 2022	0.218	-12.983	9.798		
- Soil temp: eCO <sub>2</sub>	0.449	0.655	0.828		

---

### RTD

<b>Shrub model</b>				0.74	<0.001
- Intercept	<0.001	0.285	0.056		
- Soil temp	0.355	-0.009	0.01		
- Soil moist	0.601	-0.015	0.028		
- eCO <sub>2</sub>	0.562	0.021	0.036		
- Year					

- 2015	0.155	-0.074	0.05		
- 2016	0.194	-0.124	0.092		
- 2017	-	-	-		
- 2022	0.054	0.182	0.01		
- Soil temp: eCO <sub>2</sub>	0.601	-0.004	0.008		
<b>Tree model</b>				0.33	0.054
- Intercept	<0.001	-1.605	0.173		
- Soil temp	0.144	0.05	0.033		
- Soil moist	0.112	0.148	0.088		
- eCO <sub>2</sub>	0.76	-0.037	0.118		
- Year					
- 2015	0.848	0.032	0.162		
- 2016	0.392	0.261	0.296		
- 2017	-	-	-		
- 2022*	0.016	0.771	0.285		
- Soil temp: eCO <sub>2</sub>		-0.026	0.027		

## 5. Details about statistical methods

As microtopography (hummock and hollow) did not contribute to random variation significantly in linear mixed effects models and none of the traits differed significantly between hummocks and hollows, we aggregated the root trait dataset to the plot-level by taking the mean of hummock and hollow values. We then estimated the effects of soil warming and elevated CO<sub>2</sub> treatments on the individual traits (SRL, RTD and root C:N, as well as fine root length production) using linear mixed effects models with function *lme* from package *nlme* (Pinheiro et al., 2023; Pinheiro & Bates, 2000) in R (v4.3.3 R Core Team 2024). Each model was built for trees and shrubs separately (e.g. separate tree and shrub models for SRL). In the models, soil temperature (mean of half-hourly measurements over the period of ingrowth core deployment per plot), elevated CO<sub>2</sub> treatment (ambient or elevated), soil moisture (gravimetric water content, g/g), and the interaction between soil temperature and elevated CO<sub>2</sub> treatment were fixed effects, while year was a random effect, due to sample sizes and treatment levels varying between years.

The response variables (SRL, RTD, root C:N and fine root length production) were log-transformed (except tree root C:N for which model residuals were normally distributed without transformation), after which the residuals were normally distributed in all models. To avoid multicollinearity arising from the interaction term, soil temperature and soil moisture were also centered. Multicollinearity was checked with variance inflation factor with function *vif* from package *car* (Fox & Weisberg,



2018). In tree RTD and shrub fine root length production models, year caused residual heterogeneity which was allowed for in the models by including it in *varIdent* variance structure (different variances per stratum).

We assessed the most significant trait predictors with backward variable selection using the log likelihood ratio method with AIC and *p*-values (models were fitted with maximum likelihood for model comparisons and restricted maximum likelihood for final model assessments).

To explore the effect of year on the root traits in the experiment, we used linear regressions with function *lm* from package *stats* (R Core Team, 2024), due to the lack of random effects and residual variance heterogeneity. For these models, we did not do backward variable selection but explored only the full models (table 4).

## References

Fox, J., & Weisberg, S. (2018). *An R companion to applied regression* (3rd ed.). SAGE Publications.

Pinheiro, J., & Bates, D. M. (2000). *Mixed-effects models in S and S-PLUS* (1st ed.) [PDF]. Springer.

Pinheiro, J., Bates, D. M., & R Core Team. (2023). *nlme: Linear and Nonlinear Mixed Effects Models* (Versions 3.1-164). <https://CRAN.R-project.org/package=nlme>

R Core Team. (2024). *R: A Language and Environment for Statistical Computing* (Version 4.3.3). R Foundation for Statistical Computing. <https://www.R-project.org>