

Towards Risk-Smart Restoration: Policy Insights on Disease and Ecological Risks from the RESTOREID Project



Integrating One Health into ecosystem restoration policy



Soushieta Jagadesh, Adam Fell, Marie Vandewalle, Brad Duthie, Lucinda Kirkpatrick, and Nils Bunnefeld

15.06.2026



Good afternoon!

I'll be sharing some insights from the RESTOREID project on how restoration policy currently addresses or often fails to address disease and ecological risks associated with ecosystem restoration.

Why this matters



EU Nature
Restoration
Regulation



Kunming-
Montréal Global
Biodiversity
Framework



UN decade on
Ecosystem
Restoration




Massive
Expansion of
Restoration
Projects Globally



Are they unintended consequences to restoration as landscapes
are being altered ?

We are currently witnessing an unprecedented global expansion of restoration initiatives. This is being driven by frameworks such as the EU Nature Restoration Regulation, the Kunming–Montréal Global Biodiversity Framework, and the UN Decade on Ecosystem Restoration. These initiatives are essential and urgently needed. But restoration fundamentally alters landscapes, species assemblages, and human interactions with ecosystems. And while restoration policies strongly emphasise ecological and climate benefits, far less attention has been given to possible unintended consequences. So the question is not whether to restore degraded landscapes but how to restore it intelligently.

The Double Blind Spot



The central framing of our work is what we call a ‘double blind spot’ from our systematic review of scientific literature and limited policies published early 2026.

On one side, there is a science gap.

We still have limited empirical evidence linking restoration pathways with zoonotic disease outcomes, particularly across different ecological contexts.

Disease dynamics are highly context specific, and there are relatively few long-term restoration-health monitoring systems.

On the other side, there is a policy gap.


Most restoration policies rarely request or integrate disease-related evidence, mention One Health only superficially if at all, and include very limited adaptive risk management mechanisms.

Together, these gaps reinforce one another.

Policies often assume landscape restoration is inherently positive by default.

What We Did in RESTOREID

A systematic multilingual policy appraisal of restoration and zoonotic risk integration



To explore this issue, we conducted a systematic multilingual appraisal of international and national restoration-related policies.

We reviewed policies from governments, UN agencies, WHO, FAO, IUCN, WOA, and related frameworks, using searches across four languages.

We then developed an evaluation framework examining:

restoration objectives,

zoonotic risk considerations,

interdisciplinary integration including One Health,

governance mechanisms,

and multiple dimensions of equity.

Using thematic content analysis in NVivo, we coded both explicit and implicit references to disease-related risks and compared patterns across governance scales and regions.


The goal was not to rank policies,

but to identify recurring blind spots and opportunities for improvement.

Main Findings

Restoration benefits are widely recognised, risks are largely absent!


Widely recognised in policies



Policies did not
consider
unintended risks



Largely absent from policies



Our findings revealed a very strong imbalance. Restoration policies overwhelmingly emphasise benefits such as: ecosystem services, carbon sequestration, climate resilience, and socioeconomic co-benefits. However, risks are largely absent from policy frameworks. Very few policies discussed: zoonotic spillover risk, vector dynamics, biosecurity, or social inequities linked to restoration processes.

Restoration: A Double Opportunity

Restoration can become a public health intervention when done intelligently

Challenge: Unintended Risks
may emerge



↑ Wildlife-human contact



Habitat changes ↑
reservoirs hosts/ vectors



Biosecurity risks



Social vulnerabilities

Opportunity: Regulate & reduce
disease risks



↑ Complex species
assemblages



↓ Fragmentation & ↑
ecological connectivity



↑ Ecosystem resilience



Better land governance

At this point, it's important to emphasise that restoration itself is not the problem.

What matters is how restoration is designed, implemented, and governed.

In many ways, restoration creates a double opportunity.

If ecological complexity is oversimplified, or if local context is ignored, restoration may unintentionally create conditions that favour disease emergence.

For example:

changes in habitat structure can increase wildlife-human interfaces,

certain restoration configurations may favour reservoir hosts or vectors,

biosecurity risks can emerge through planting materials or species translocations,


and restoration processes can sometimes deepen social vulnerabilities or land-use tensions. many of the social risks associated with restoration are not ecological problems alone but governance challenges. But the opposite is also true. When restoration strengthens ecological integrity, it can help regulate disease dynamics rather than amplify them.

Restoring more complex species assemblages, improving ecological connectivity, reducing ecosystem degradation, and strengthening governance systems by explicitly including equity, and participation can all contribute to more resilient landscapes and healthier human-environment relationships. So restoration should not simply be viewed as a biodiversity intervention. It can also become a form of preventative public health infrastructure.

And that shift in perspective is critical if restoration is going to remain effective and socially legitimate in the long term.

The 3 pillars of “Risk-Smart” Restoration

From “Risky Restoration” to “Risk-Smart Restoration”



So what would ‘risk-smart restoration’ actually look like?

Our framework identifies three core pillars.

The first is moving from generic restoration to context-aware design.

That means thinking not only about planting trees or restoring habitat, but about which species assemblages are being promoted, how landscapes are connected, and where high-risk wildlife-human interfaces may emerge.

In other words, restoration design itself becomes part of preventative health planning.

The second shift is from static restoration targets to adaptive monitoring.


Most policies monitor hectares restored or carbon captured but far fewer monitor ecological imbalance, vector dynamics, or potential spillover pathways.

A risk-smart approach treats restoration as an ongoing learning process rather than a one-time intervention.

And the third shift is from siloed governance to inclusive and integrated governance.

Health, biodiversity, agriculture, and land-use planning are still often treated as separate policy domains. Yet zoonotic risk emerge at the intersection of these systems. Effective restoration governance therefore requires more than cross-sector collaboration. It also requires attention to who benefits from restoration, who bears the costs, and whose voices are included in decision-making.

Integrating public health expertise, ensuring equitable benefit-sharing, incorporating social safeguards, and engaging local communities meaningfully can help reduce vulnerabilities while strengthening the legitimacy and long-term success of restoration initiatives. Together, these three pillars help move restoration from a potentially reactive process toward something far more resilient, adaptive, and socially sustainable.



So how do we translate these ideas into actual policy and implementation?

What we propose is not a separate health framework layered onto restoration but a restoration lifecycle that integrates health considerations from beginning to end.

That starts before restoration even begins.

At the planning stage, policies should move beyond generic restoration targets and ask more context-specific questions:

What ecosystems are being restored?

Which species or vectors may respond to these landscape changes?

Where might new wildlife-human interfaces emerge?

In other words, risk anticipation needs to become part of restoration design itself.

Then during implementation, restoration should function as an adaptive process.

Ecosystems are dynamic, and restoration outcomes rarely unfold exactly as predicted.

So monitoring systems, stakeholder engagement, and data sharing become essential not only for biodiversity outcomes, but also for identifying unintended ecological or health signals early enough to respond.


And finally, restoration does not end once trees are planted or targets are achieved.

Long-term surveillance, transparent governance, and early-warning systems are critical for ensuring that restored landscapes remain resilient, socially legitimate, and safe over time.

Conclusion



One Health
strengthens
restoration
legitimacy &
resilience



One-size-fits-
all restoration
can backfire



Risk-smart
Restoration is
the way
forward

I'd like to conclude with three key messages.

First: integrating One Health principles strengthens restoration legitimacy, resilience, and long-term sustainability

Second: one-size-fits-all restoration approaches can backfire if local ecological and social contexts are ignored.

And Ultimately, risk smart restoration is a powerful opportunity for biodiversity and health co-benefits provided risks are anticipated and managed from the outset.

The question is not whether to restore nature, but how to restore it intelligently.

Thank you.

Thank you

Restoration is a powerful opportunity for biodiversity and health co-benefits, provided One Health is embedded throughout the restoration lifecycle.



Policy brief

Reducing zoonotic risk through nature restoration

Context, challenges and recommendations for National Restoration Plans in the implementation of the Nature Restoration Regulation.

Nature restoration and health

The EU Nature Restoration Regulation (NRR)¹, adopted in August 2024, sets ambitious targets to restore degraded ecosystems by 2030 and beyond. Research has, however, identified risks: **over 75% of emerging infectious diseases are zoonotic**, and land-use change is a major driver of spillover risk. Restoration is a primary tool to reverse the land-use changes that drive spillover events, moving from unstable, simplified ecosystems to resilient, self-regulating ones.

The key policy question is **how to ensure restoration strengthens ecosystem resilience while mitigating existing zoonotic disease risks**.



Key messages

- **Nature restoration is a proactive public health strategy.** Intact and biodiverse ecosystems act as a “biological buffer,” stabilising pathogen dynamics and reducing the long-term risk of zoonotic spillover compared to degraded landscapes.
- **Zoonoses are driven by habitat destruction.** While degradation creates the primary risk, well-planned restoration provides the solution by rebuilding ecological complexity. High-quality restoration reduces the dominance of high-risk “reservoir” species (like certain rodents) through the “dilution effect.”
- **Risk management ensures success.** While restoration is a net positive for health, “safe-by-design” planning is essential to manage localised, context-specific risks during the transition phase, ensuring public trust and social acceptability.
- The **One Health** approach, when integrated from the design stage of National Restoration Plans, enables policymakers **to anticipate, reduce and manage zoonotic risks without compromising ecological or health benefits**.

1. Footnotes can be found in the information sheet online.





Key observations

Understanding the zoonotic risks associated with nature restoration

What is a “zoonotic risk”?

A zoonotic risk is the **probability that a pathogen of animal origin will be transmitted to humans**, leading to infection, with health, social, and economic consequences. This risk depends on three interacting components:

- animal reservoirs (wildlife or domestic fauna),
- pathogens (e.g., viruses, bacteria, protozoa) and vectors (e.g., ticks, mosquitoes),
- intensity of human–wildlife contact.

The highest risk currently exists in fragmented, degraded edge habitats. Nature restoration can either reduce or amplify these risks, depending on the design and implementation of the restoration project.

How restoration can reduce or amplify zoonotic risks?

The relationship between land degradation, ecosystem complexity (compositional, structural, functional), and zoonotic spillover is context-dependent, varying with ecosystem type, pathogen ecology, species involved, and patterns of human use. However, we present general key points which should be considered when planning and implementing restoration measures.

a. Modification of animal communities

Habitat restoration changes wildlife populations in space and time. In low-diversity ecosystems, a few highly competent reservoir species (species that are effective at hosting, maintaining and transmitting pathogens to vectors e.g., certain rodents or deer supporting tick cycles) may dominate. By contrast, **complex species assemblages**, meaning diverse, well-balanced communities that include a diverse array of predators, competitors and multiple host species, tend to reduce dominance of high-risk hosts and stabilise disease dynamics (the “dilution effect”).

b. Change in vector abundance

Wetland, forest, or landscape restoration can modify vector habitats and human access patterns. When well designed, with appropriate habitat structure, connectivity, and trophic balance, restoration strengthens biodiversity and helps regulate host–vector systems rather than amplifying risk.

c. Human-wildlife interactions

Restoration projects are often accompanied by new recreational or ecotourism activities, increasing contact between humans, wildlife, and vectors. As ecosystems require time to stabilise, including reaching balanced host–pathogen dynamics, **recreational access should be phased and aligned with ecological recovery timelines to minimise exposure risks**. However, the risk of inaction outweighs the manageable transition risks of restoration.

Handling risks at the governing level

Political and social risks

A systematic appraisal of 30 national and international policies on landscape restoration, zoonoses, and related health frameworks published since 2015 found that while most highlight ecological and socio-economic benefits, **only 10% address unintended health or social consequences**. Health sectors are rarely involved, and monitoring overlooks these aspects. This lack of anticipation can fuel controversy, erode trust, and politicise health risks to oppose restoration.

Integrating zoonotic risk through a One Health approach

The **One Health approach** acknowledges the interdependence of human, animal and ecosystem health, making it particularly relevant to restoration projects that reshape habitats, wildlife, and land use. Integrating One Health considerations into national restoration plans would:

- **Strengthen policy coherence** by involving health and veterinary authorities from the design stage;
- **Mobilise joint expertise** in ecology and epidemiology;
- **Improve data sharing** between environmental and health sectors.

Restore nature and reduce zoonotic risks

Landscape degradation (e.g., deforestation, fragmentation, biodiversity loss) is consistently linked to increased disease risk (red), especially for mosquito-, bat- and rodent-borne diseases. In contrast, several

restoration actions are associated with reduced or neutral effects (blue or white). These observations are gathered in the “heat map” shown in Figure 1.




Figure 1: Heat map showing the size and direction of reported effects of land-use changes (LUC) and biodiversity–disease relationships on disease transmission risk across different disease groups (vector-borne diseases in blue, zoonoses in pink, general in gold, neglected diseases in grey, wildlife diseases in green and tuberculosis in purple): blue with a downward arrow indicates a decreased disease risk, red with an upward arrow indicates an increased risk and white with a bidirectional arrow indicates a reported non-significant effect.

Well-designed restoration reduces zoonotic risks by:

- Rebuilding complex species assemblages**, i.e., diverse, balanced communities, that prevent dominance of highly competent reservoir hosts;
 - Improving habitat quality and connectivity**, reducing ecological stress and unstable host–pathogen dynamics;
 - Avoiding fragmented, simplified landscapes**, which the heat map shows are frequently associated with increased transmission risk.
- In short, restoration reduces risks already generated by ecological degradation when it strengthens ecological (i.e., functional and structural) complexity and stability.




Figure 2: When nature restoration reduces disease risk and when it doesn't, www.restoreid.eu



Recommendations for national policy makers

1. Design (Conception and Planning)

Restoration design should integrate context-specific zoonotic risk considerations from the outset (Articles 4–13), supported by the Context-Based Decision Pathway¹.

- Apply a “safe by design” approach prioritising restoration that promotes trophic recovery and complex species assemblages, reducing dominance of highly competent reservoir hosts.
- Undertake proportionate ex ante screening to identify:
 - key species assemblages,
 - potential vectors and reservoir hosts,
 - human–wildlife interface zones.
- Design for local context by adapting restoration to ecosystem type², land-use history, fragmentation, and endemic disease landscapes.
- Manage interfaces and access:
 - reduce high-risk edge habitats,
 - regulate access to restored areas, particularly during early recovery phases.

See Box 2³ for specificities of each ecosystem targeted by NRR articles.

2. Monitoring

Monitoring should enable adaptive management of restoration outcomes under Articles 20–21, ensuring biodiversity gains are achieved while maintaining low zoonotic risk.

- Integrate One Health indicators into monitoring systems, including:
 - species assemblage composition,
 - host or vector presence where relevant,
 - ecosystem resilience indicators.
- Apply context-specific monitoring, tailored to ecosystem type and risk drivers (no one-size-fits-all approach).
- Use existing surveillance systems to contextualise restoration:
 - national systems and the European Centre for Disease Prevention and Control for vector and disease trends.
- Leverage existing data platforms:
 - Global Biodiversity Information Facility and European Environment Agency for biodiversity monitoring,
 - MOOD platform for integrating environmental and health data.

Monitoring should support early detection of ecological imbalance or zoonotic emergence signals and inform timely design adjustments⁴.

1. Box 1, see information sheet online.

2. Checklists 1, 2 and 3, see information sheet online.

3. Box 2, see information sheet online.

4. Table 1 (indicators to monitor zoonotic risks), see information sheet online.



3. Governance and Implementation

Effective implementation requires clear roles, cross-sector coordination, and transparent processes aligned with NRR provisions (Articles 14–19).

- Strengthen transboundary coordination (Article 14) for shared ecosystems through defined authorities and data-sharing arrangements.
- Integrate health into governance involving public health and veterinary authorities in National Restoration Plans (NRPs).
- Operationalise stakeholder engagement by establishing consultation, grievance, and benefit-sharing mechanisms.
- Build capacity to support cross-sector training and exchanges (environment, agriculture, health, planning).
- Ensure data transparency and usability:
 - provide access to data used in planning and monitoring,
 - identify responsible authorities and support interpretation.
- Communicate clearly on both the benefits of restoration and the management of human–wildlife interactions to support social acceptance and avoid conflicts.



Conclusion: Nature restoration as a lever for One Health

The NRR offers a pivotal opportunity to reshape our relationship within nature. In an era of emerging infectious diseases, human health cannot remain peripheral to environmental policy. Integrating zoonotic risk through a One Health approach does not weaken restoration ambition but strengthens its credibility, resilience and public legitimacy. This integration would allow restoration to become, not just an ecological imperative, but a public health strategy.

The question is not whether to restore nature, but how to restore it intelligently, recognising that ecological integrity and human health are inseparable.

Nature restoration can reduce disease risk – but only if it is designed with health in mind. One-size-fits-all can backfire.



Link to sources

BiodivRestore Knowledge Hub

The scientific publications used in this policy brief can be found in the information sheet of this briefing, downloadable from: www.biodiversa.eu/policy-briefs/

Photos: Unsplash, Pixabay.

Contact

contact@biodiversa.eu
www.biodiversa.eu



@Biodiversa.eu



@BiodiversaPlus

About this Policy brief

This Policy Brief is part of a series aiming to inform policymakers involved in the implementation of the [Nature Restoration Regulation](#) with policy recommendations based on the expertise of the BiodivRestore Knowledge Hub experts.

The series of Biodiversa+ Policy briefs can be found at www.biodiversa.eu/policy-briefs/.

This publication emerged from the BiodivRestore Knowledge Hub, was funded and supervised by Biodiversa+, and produced by Robin Goffaux, Cloé Durieux and Julie de Bouville.

The key research results presented were co-drafted and validated by a researcher from the BiodivRestore Knowledge Hub: Dr Lucinda Kirkpatrick, as well as external contributors: Dr Soushieta Jagadesh, Prof. Nadja Kabisch, Prof. Frauke Ecke, Prof. Nils Bunnefeld.

With thanks to Bastian Bertschy, Marlies Laethem, Dr. Ron Winkler, Dr. Magnus Tannerfeldt, Luciana Carotenuto, Dr. Giulia Albani Rocchetti.

Funded by the European Union. Views and opinions expressed are, however, those of the authors only and do not necessarily reflect those of the European Union or the granting authority, the Research Executive Agency (REA). Neither the European Union nor the granting authority can be held responsible for them.



Co-funded by
the European Union
under Grant Agreement
No 642420



Produced in May 2026.